

***K*-THEORY OF NON-ARCHIMEDEAN RINGS II**

MORITZ KERZ^{id}, SHUJI SAITO^{id} AND GEORG TAMME^{id}

Abstract. We study fundamental properties of analytic K -theory of Tate rings such as homotopy invariance, Bass fundamental theorem, Milnor excision, and descent for admissible coverings.

§1. Introduction

This note is a sequel to our note [7], and we refer to that paper for a general introduction and motivation. In part I, we were mainly concerned with the construction of what we call analytic K -theory of Tate rings and a comparison with continuous K -theory as proposed by Morrow [12]. The main result there was a weak equivalence between continuous and analytic K -theory for regular Noetherian Tate rings admitting a ring of definition satisfying a certain weak resolution of singularities property. Negative continuous K -theory was studied by Dahlhausen in [1].

The goal here is to study more closely the properties of analytic K -theory. It turns out that analytic K -theory has good formal properties, similar to Weibel’s homotopy K -theory, without any regularity assumption. However, we have to restrict to Tate rings admitting a Noetherian, finite-dimensional ring of definition. In particular, for such Tate rings, we establish the analytic Bass fundamental theorem (Corollary 2.6), pro-homotopy invariance (Corollary 2.7), Milnor excision (Theorem 3.2), and descent for rational coverings of the associated adic space (Theorem 4.1). The latter allows us to globalize the definition of analytic K -theory to adic spaces satisfying a finiteness condition. In addition, we use our results to prove that analytic K -theory of such Tate rings is in fact weakly equivalent to the “ \mathbb{A}^1 -localization” of Morrow’s continuous K -theory (see [12]; see also Corollary 2.9). We thank the referee for suggesting to do so.

Our main tool is a fiber sequence relating the continuous K -theory of a ring of definition, or more generally a Raynaud type model, with the analytic K -theory of a Tate ring (Theorems 2.4 and 2.10).

1.1 Notation and conventions

By a π -adic (or just adic) ring, we mean a topological ring A_0 whose topology is the π -adic one for an element $\pi \in A_0$. We usually assume that A_0 is complete. A topological ring A is called a Tate ring if it admits an open subring $A_0 \subseteq A$ and an element $\pi \in A_0$ such that the subspace topology on A_0 is the π -adic one, and π is invertible in A . Such an A_0 is called a ring of definition, and π is called a pseudo-uniformizer. We have $A = A_0[\frac{1}{\pi}]$. If A is Tate, then a subring $A_0 \subseteq A$ is a ring of definition if and only if it is open and bounded. The basic reference is [4, §1].

Received May 15, 2022. Revised January 8, 2023. Accepted January 28, 2023.
2020 Mathematics subject classification: 19D25, 14G22.

Kerz was supported by the Deutsche Forschungsgemeinschaft (DFG) through CRC 1085. Saito was supported by JSPS KAKENHI Grant (15H03606). Tamme was supported by the Deutsche Forschungsgemeinschaft (DFG) through CRC 1085 and TRR 326 (Project-ID 444845124).

© The Author(s), 2023. Published by Cambridge University Press on behalf of Foundation Nagoya Mathematical Journal. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

We denote by \mathbf{Spc} and \mathbf{Sp} the ∞ -categories of spaces and spectra, respectively. $\mathrm{Pro}(\mathbf{Sp})$ denotes the ∞ -category of pro-spectra (see, e.g., [7, §2]). A map of pro-spectra f is a weak equivalence, if its level-wise truncation $\tau_{\leq n}f$ is an equivalence of pro-spectra for every integer n (see [7, §2.2] for a discussion and equivalent characterizations). Similarly, a sequence $X \rightarrow Y \rightarrow Z$ of pro-spectra is a weak fiber sequence if it is weakly equivalent to a fiber sequence.

The letter K denotes the nonconnective algebraic K -theory functor for rings and quasi-compact, quasi-separated schemes. Its connective covering is denoted by k . For the following notations, we refer to the corresponding definitions in [7]:

- K^{cont} for an adic ring: Definition 5.1.
- K^{cont} for a Tate ring: Definition 5.3.
- k^{an} for an adic ring: Definition 6.5.
- k^{an} for a Tate ring: Definition 6.11. (Warning: in general, this is not the connective cover of K^{an} .)
- K^{an} for a Tate ring: Definition 6.15.

§2. Recollections from part I, and complements

2.1 Pro-homotopy algebraic K -theory

In this subsection, we introduce a variant of Weibel's homotopy algebraic K -theory, which will be an important tool to compare the analytic K -theory of a Tate ring with the continuous K -theory of a ring of definition without assuming any regularity condition (see [7, §6.1] for similar constructions in the setting of complete Tate rings).

Let R be a ring, and let $\pi \in R$ be any element. For nonnegative integers p, j , we define

$$R[\Delta_{\pi^j}^p] := R[t_0, \dots, t_p]/(t_0 + \dots + t_p - \pi^j).$$

For fixed j and varying p , the $R[\Delta_{\pi^j}^p]$ form a simplicial ring, which we denote by $R[\Delta_{\pi^j}]$. Multiplication of the coordinates t_i by π induces maps of simplicial rings $R[\Delta_{\pi^j}] \rightarrow R[\Delta_{\pi^{j-1}}]$, and in this way we obtain the pro-simplicial ring

$$\text{“}\lim_j\text{” } R[\Delta_{\pi^j}].$$

Let \mathbf{Sch}_R denote the category of quasi-compact, quasi-separated (qcqs) R -schemes. Let F be a functor $F: \mathbf{Sch}_R^{\mathrm{op}} \rightarrow \mathbf{Sp}$. We define a new functor $F^{\pi^\infty}: \mathbf{Sch}_R^{\mathrm{op}} \rightarrow \mathrm{Pro}(\mathbf{Sp})$ as follows. We let $F^{\pi^j}(X)$ be the geometric realization of the simplicial spectrum $[p] \mapsto F(X \otimes_R R[\Delta_{\pi^j}^p])$ and define

$$F^{\pi^\infty}(X) = \text{“}\lim_j\text{” } F^{\pi^j}(X) \in \mathrm{Pro}(\mathbf{Sp}).$$

DEFINITION 2.1. Let X be a qcqs R -scheme. We define its *pro-homotopy algebraic K -theory (with respect to $\pi \in R$)* as the pro-spectrum

$$K^{\pi^\infty}(X).$$

Similarly we define a version with support on $X/(\pi) = X \otimes_R R/(\pi)$:

$$K^{\pi^\infty}(X \text{ on } (\pi)).$$

For an R -scheme X , we write $X[t] = X \otimes_R R[t]$. By construction, K^{π^∞} satisfies the following form of pro-homotopy invariance.

LEMMA 2.2. For $X \in \mathbf{Sch}_R$, the canonical map

$$K^{\pi^\infty}(X) \rightarrow \text{“lim”}_{t \rightarrow \pi t} K^{\pi^\infty}(X[t])$$

is an equivalence of pro-spectra. Similarly for $K^{\pi^\infty}(-\text{on}(\pi))$.

The proof is the same as that of [7, Prop. 6.3]. We will use the following easy observation.

LEMMA 2.3. If π is invertible on X , there is a natural equivalence

$$K^{\pi^\infty}(X) \cong KH(X),$$

where $KH(X)$ denotes Weibel’s homotopy K -theory spectrum. In particular, for any X , there is a fiber sequence of pro-spectra

$$K^{\pi^\infty}(X \text{ on}(\pi)) \rightarrow K^{\pi^\infty}(X) \rightarrow KH(X[\frac{1}{\pi}]).$$

Proof. If π is invertible on X , then the transition map $R[\Delta_{\pi^j}^p] \rightarrow R[\Delta_{\pi^0}^p] = R[\Delta^p]$ induces an isomorphism $X \otimes_R R[\Delta^p] \rightarrow X \otimes_R R[\Delta_{\pi^j}^p]$. Thus, the pro-system defining $K^{\pi^\infty}(X)$ is constant with value $KH(X)$. The second claim follows from the first one together with the localization sequence associated with the identity map $X \rightarrow X[\frac{1}{\pi}]$. \square

2.2 The fundamental fiber sequence

The following weak fiber sequence and its generalization to non-affine models in Theorem 2.10 are our fundamental tool to control the analytic K -theory of Tate rings. For a π -adic ring A_0 , we denote by $K^{\text{cont}}(A_0)$ the pro-spectrum “lim” $_n K(A_0/(\pi^n))$.

THEOREM 2.4. Let A_0 be a complete π -adic ring, and let $A = A_0[1/\pi]$ be the associated Tate ring. Assume that A_0 is Noetherian and of finite Krull dimension. Then there is a weak fiber sequence of pro-spectra

$$K^{\pi^\infty}(A_0 \text{ on}(\pi)) \rightarrow K^{\text{cont}}(A_0) \rightarrow K^{\text{an}}(A).$$

Before entering the proof, we recall the analytic Bass construction, referring to [7, §4.4] for details. For a functor E defined on the category of adic or Tate rings and with values in a cocomplete stable ∞ -category (such as \mathbf{Sp} or $\text{Pro}(\mathbf{Sp})$), we define the functor ΛE by

$$\Lambda E(A) = \text{fib}(E(A\langle t \rangle) \sqcup_{E(A)} E(A\langle t^{-1} \rangle) \rightarrow E(A\langle t, t^{-1} \rangle)).$$

If E has the structure of a module over connective algebraic K -theory, then there is a natural map $\lambda: E(A) \rightarrow \Lambda E(A)$ given as the composite

$$\lambda: E(A) \xrightarrow{\sim t} \Omega E(A\langle t, t^{-1} \rangle) \rightarrow \Lambda E(A).$$

If λ is an equivalence, then E satisfies an analytic version of the Bass fundamental theorem (see [7, Prop. 4.13]). The analytic Bass construction E^B of E is then given by

$$E^B(A) = \text{colim}(E(A) \xrightarrow{\lambda} \Lambda E(A) \xrightarrow{\Lambda(\lambda)} \Lambda^2 E(A) \rightarrow \dots).$$

EXAMPLE 2.5. Let k^{cont} be the functor from adic rings to pro-spectra that assigns to A_0 the connective continuous K -theory

$$k^{\text{cont}}(A_0) = \text{“lim”}_n k(A_0/(\pi^n)).$$

As a rule, we write k for the connective algebraic K -theory $\tau_{\geq 0}K$. Then $k^{\text{cont},B}$ is canonically equivalent to nonconnective continuous K -theory K^{cont} . Indeed, using the isomorphisms $A_0\langle t \rangle/(\pi^n) \cong A_0/(\pi^n)[t]$ and similarly for t and t^{-1} , the classical Bass fundamental theorem [14, Th. 6.6] implies that $\lambda: K^{\text{cont}} \rightarrow \Lambda K^{\text{cont}}$ is an equivalence, hence $K^{\text{cont}} \simeq K^{\text{cont},B}$. Moreover, by nil-invariance of negative K -groups, the fiber of the canonical map $k^{\text{cont}} \rightarrow K^{\text{cont}}$ has values in constant, coconnective pro-spectra. Hence, this fiber vanishes upon applying the analytic Bass construction (see [7, Lem. 4.15]).

Proof of Theorem 2.4. As in [7, §3.1], we denote by $A_0\langle \Delta_{\pi^j}^p \rangle$ the π -adic completion of the ring $A_0[\Delta_{\pi^j}^p]$. In [7, Def. 6.5], we defined the pro-spectrum

$$k^{\text{an}}(A_0) = \text{“lim”}_j k(A_0\langle \Delta_{\pi^j} \rangle),$$

where $k(A_0\langle \Delta_{\pi^j} \rangle)$ is the geometric realization of the simplicial spectrum $[p] \mapsto k(A_0\langle \Delta_{\pi^j}^p \rangle)$. Similarly, we set¹

$$k^{\text{an}}(A_0 \text{ on } (\pi)) = \text{“lim”}_j k(A_0\langle \Delta_{\pi^j} \rangle \text{ on } (\pi)),$$

where $k(- \text{ on } (\pi))$ is the connective covering of $K(- \text{ on } (\pi))$. There is a natural map $k^{\text{an}}(A_0 \text{ on } (\pi)) \rightarrow k^{\text{an}}(A_0)$ and the pro-spectrum $\tilde{k}^{\text{an}}(A; A_0)$ is defined to fit in the cofiber sequence

$$k^{\text{an}}(A_0 \text{ on } (\pi)) \rightarrow k^{\text{an}}(A_0) \rightarrow \tilde{k}^{\text{an}}(A; A_0).$$

By [7, Th. 6.9], we have a natural weak equivalence $k^{\text{an}}(A_0) \simeq k^{\text{cont}}(A_0)$, and thus we have a natural weak fiber sequence

$$k^{\text{an}}(A_0 \text{ on } (\pi)) \rightarrow k^{\text{cont}}(A_0) \rightarrow \tilde{k}^{\text{an}}(A; A_0).$$

We claim that applying the analytic Bass construction $(-)^B$ yields the asserted weak fiber sequence. Note that $(-)^B$ preserves weak equivalences and weak fiber sequences of functors as its construction only involves colimits.

First, by [7, Lem. 6.18(i)], there is a natural weak equivalence $(\tilde{k}^{\text{an}})^B(A; A_0) \simeq K^{\text{an}}(A)$. Second, by Example 2.5, there is an equivalence $k^{\text{cont},B}(A_0) \cong K^{\text{cont}}(A_0)$. It thus remains to prove that the analytic Bass construction of $k^{\text{an}}(- \text{ on } (\pi))$ evaluated on A_0 is weakly equivalent to $K^{\pi^\infty}(A_0 \text{ on } (\pi))$.

Note that by Thomason’s excision (see [14, Prop. 3.19]), we have equivalences

$$\begin{aligned} K(A_0\langle \Delta_{\pi^j}^p \rangle \text{ on } (\pi)) &\simeq K(A_0[\Delta_{\pi^j}^p] \text{ on } (\pi)), \\ K(A_0\langle \Delta_{\pi^j}^p \rangle \langle t \rangle \text{ on } (\pi)) &\simeq K(A_0[\Delta_{\pi^j}^p][t] \text{ on } (\pi)), \end{aligned}$$

and similarly for t^{-1} and t, t^{-1} . By Weibel’s K -dimension conjecture, proven in [8], all these spectra are $(-d-1)$ -connective for $d = \dim(A_0)$. The classical Bass fundamental theorem and excision thus imply that

¹ In [7], this pro-spectrum is denoted $\bar{k}^{\text{an}}(A_0)$.

$$\Lambda^n k(A_0 \langle \Delta_{\pi_j}^p \rangle \text{ on } (\pi)) \simeq K(A_0 \langle \Delta_{\pi_j}^p \rangle \text{ on } (\pi)) \simeq K(A_0 [\Delta_{\pi_j}^p] \text{ on } (\pi))$$

for all p and j and every $n > d$. As finite limits and finite colimits in $\text{Pro}(\mathbf{Sp})$ are computed level wise and as geometric realizations in \mathbf{Sp} commute with finite limits and colimits this shows that

$$\Lambda^n k^{\text{an}}(A_0 \text{ on } (\pi)) \simeq \Lambda^n K^{\pi^\infty}(A_0 \text{ on } (\pi)) \simeq K^{\pi^\infty}(A_0 \text{ on } (\pi))$$

for $n > d$. So the diagram in the definition of the analytic Bass construction for $E = k^{\text{an}}(- \text{ on } (\pi))$, evaluated on A_0 , is eventually constant with value $K^{\pi^\infty}(A_0 \text{ on } (\pi))$. This concludes the proof. \square

We record the following applications.

COROLLARY 2.6. (*Bass fundamental theorem for analytic K-theory*) *Let A be a complete Tate ring which admits a finite-dimensional, Noetherian ring of definition. Then the map $\lambda: K^{\text{an}}(A) \rightarrow \Lambda K^{\text{an}}(A)$ is a weak equivalence of pro-spectra, and hence the Bass fundamental theorem holds. That is, for every integer i , we have an exact sequence*

$$0 \rightarrow K_i^{\text{an}}(A) \rightarrow K_i^{\text{an}}(A\langle t \rangle) \oplus K_i^{\text{an}}(A\langle t^{-1} \rangle) \rightarrow K_i^{\text{an}}(A\langle t, t^{-1} \rangle) \rightarrow K_{i-1}^{\text{an}}(A) \rightarrow 0$$

and the right-hand map has a splitting given by multiplication with the class of $t \in K_1(A\langle t, t^{-1} \rangle)$.

Proof. Let A_0 be a finite-dimensional Noetherian ring of definition. As in Example 2.5, the map $\lambda: K^{\text{cont}}(A_0) \rightarrow \Lambda K^{\text{cont}}(A_0)$ is an equivalence. Using again Thomason’s excision and the classical Bass fundamental theorem, we see that also $\lambda: K^{\pi^\infty}(A_0 \text{ on } (\pi)) \rightarrow \Lambda K^{\pi^\infty}(A_0 \text{ on } (\pi))$ is an equivalence. By Theorem 2.4, this implies that $\lambda: K^{\text{an}}(A) \rightarrow \Lambda K^{\text{an}}(A)$ is a weak equivalence. The rest now follows from [7, Prop. 4.13]. \square

COROLLARY 2.7. (*Pro-homotopy invariance*) *Let A be a complete Tate ring which admits a finite-dimensional, Noetherian ring of definition. Then the map*

$$K^{\text{an}}(A) \rightarrow \text{“}\lim_{t \rightarrow \pi t}\text{” } K^{\text{an}}(A\langle t \rangle)$$

is a weak equivalence.

Proof. Both other terms in the fundamental fiber sequence of Theorem 2.4 satisfy this type of pro-homotopy invariance (see [7, Lem. 5.13] for K^{cont} and Lemmas 2.2 and 2.3 for $K^{\pi^\infty}(- \text{ on } (\pi))$). \square

We want to compare analytic K -theory of Tate rings to the “ \mathbb{A}^1 -localization” of Morrow’s continuous K -theory [12] (see [7, Def. 5.3]), similarly as Weibel’s homotopy K -theory is the \mathbb{A}^1 -localization of algebraic K -theory. Recall that the ∞ -category $\text{Pro}(\mathbf{Sp})$ is cocomplete, so it admits geometric realizations of simplicial objects. Note also that general colimits in $\text{Pro}(\mathbf{Sp})$ cannot be computed level wise. However, up to weak equivalence, this is the case for geometric realizations of bounded below simplicial objects, as we prove in the following lemma. Let $X_\bullet = \text{“}\lim_\lambda X_\bullet^{(\lambda)}\text{”}$ be a pro-object of simplicial spectra, that is, an object of $\text{Pro}(\text{Fun}(\Delta^{\text{op}}, \mathbf{Sp}))$. We can view X_\bullet as a simplicial object in $\text{Pro}(\mathbf{Sp})$ and form its geometric realization $|X_\bullet|$ there, or we can form the level-wise geometric realizations $|X_\bullet^{(\lambda)}|$ in spectra and consider the pro-spectrum “ $\lim_\lambda |X_\bullet^{(\lambda)}|$ ”.

LEMMA 2.8. *In the above situation, assume that X_\bullet is uniformly bounded below, that is, that there exists an integer N such that $X_p^{(\lambda)} \in \mathbf{Sp}_{\geq N}$ for all $p \in \Delta$ and all $\lambda \in \Lambda$. Then the evident map*

$$|X_\bullet| \rightarrow \text{“}\lim_{\lambda}\text{” } |X_\bullet^{(\lambda)}|$$

is a weak equivalence.

Proof. It is easy to check that the ∞ -category $\text{Pro}(\mathbf{Sp})$ admits a t -structure whose connective part $\text{Pro}(\mathbf{Sp})_{\geq 0}$ is given by the essential image of the evident functor $\text{Pro}(\mathbf{Sp}_{\geq 0}) \rightarrow \text{Pro}(\mathbf{Sp})$ and whose coconnective part $\text{Pro}(\mathbf{Sp})_{\leq 0}$ is the essential image of the functor $\text{Pro}(\mathbf{Sp}_{\leq 0}) \rightarrow \text{Pro}(\mathbf{Sp})$. Its heart is the category of pro-abelian groups $\text{Pro}(\mathbf{Ab})$. The homotopy group functors of the t -structure $\pi_n: \text{Pro}(\mathbf{Sp}) \rightarrow \text{Pro}(\mathbf{Ab})$ are just the level-wise homotopy group functors $\text{Pro}(\pi_n)$, which we previously also denoted by π_n .

By shifting X_\bullet , we may assume that $N = 0$ so that X_\bullet determines a simplicial object of $\text{Pro}(\mathbf{Sp})_{\geq 0}$. In this situation, there is a convergent first quadrant spectral sequence of pro-abelian groups $E_{p,q}^1 \Rightarrow \pi_{p+q}(|X_\bullet|)$ where $E_{*,q}^1$ is the normalized chain complex associated with the simplicial pro-abelian group $\pi_q(X_\bullet)$ (see [11, Prop. 1.2.4.5]). On the other hand, for each λ , we have a convergent spectral sequence of abelian groups $E_{p,q}^{1,(\lambda)} \Rightarrow \pi_{p+q}(|X_\bullet^{(\lambda)}|)$ where again the E^1 -terms are given by the normalized chain complexes associated with the simplicial abelian group $\pi_q(X_\bullet^{(\lambda)})$. As finite limits and finite colimits in $\text{Pro}(\mathbf{Ab})$ are computed level wise, the pro-system of these spectral sequences yields a convergent spectral sequence “ \lim_{λ} ” $E_{p,q}^{1,(\lambda)} \Rightarrow$ “ \lim_{λ} ” $\pi_{p+q}(|X_\bullet^{(\lambda)}|)$. By the same reason, and as $\pi_q(X_p) = \text{“}\lim_{\lambda}\text{” } \pi_q(X_p^{(\lambda)})$, the canonical map of spectral sequences $E_{*,*}^* \rightarrow \text{“}\lim_{\lambda}\text{” } E_{*,*}^{*,(\lambda)}$ is an isomorphism on E^1 -pages. Hence, $\pi_n(|X_\bullet|) \rightarrow \text{“}\lim_{\lambda}\text{” } \pi_n(|X_\bullet^{(\lambda)}|)$ is an isomorphism of pro-abelian groups, as was to be shown. \square

Now, let A be a Tate ring with ring of definition A_0 and pseudo-uniformizer π . Recall that the continuous K -theory of A with respect to the ring of definition A_0 is given by

$$K^{\text{cont}}(A; A_0) = \text{cofib}(K(A_0 \text{ on } (\pi)) \rightarrow K^{\text{cont}}(A_0)),$$

and this pro-spectrum is independent of the choice of A_0 up to weak equivalence (see [7, Def. 5.3 and Prop. 5.4]). As this definition is clearly functorial in the pair (A_0, π) , we get a well-defined simplicial pro-spectrum $K^{\text{cont}}(A\langle\Delta_{\pi^j}^\bullet\rangle; A_0\langle\Delta_{\pi^j}^\bullet\rangle)$ (which can also be realized as a pro-simplicial spectrum) and we define $K^{\text{cont}}(A\langle\Delta_{\pi^j}\rangle; A_0\langle\Delta_{\pi^j}\rangle)$ to be its geometric realization. As weak equivalences are preserved under colimits, this definition does not depend on the choice of A_0 up to weak equivalence. Taking the colimit over all rings of definition A_0 gives the well-defined pro-spectrum $K^{\text{cont}}(A\langle\Delta_{\pi^j}\rangle)$. The “ \mathbb{A}^1 -localization” of continuous K -theory at the Tate ring A is then the pro-spectrum

$$\text{“}\lim_j\text{” } K^{\text{cont}}(A\langle\Delta_{\pi^j}\rangle).$$

COROLLARY 2.9. (\mathbb{A}^1 -localization) *Suppose the Tate ring A admits a Noetherian, finite-dimensional ring of definition. Then there is a canonical weak equivalence of pro-spectra*

$$\text{“}\lim_j\text{” } K^{\text{cont}}(A\langle\Delta_{\pi^j}\rangle) \simeq K^{\text{an}}(A).$$

Proof. Let $A_0 \subseteq A$ be a Noetherian, finite-dimensional ring of definition, and let $\pi \in A_0$ be a pseudo-uniformizer. By the discussion above, we have a weak fiber sequence of

pro-simplicial spectra

$$K(A_0\langle\Delta_{\pi^j}^\bullet\rangle \text{ on } (\pi)) \rightarrow K^{\text{cont}}(A_0\langle\Delta_{\pi^j}^\bullet\rangle) \rightarrow K^{\text{cont}}(A\langle\Delta_{\pi^j}^\bullet\rangle; A_0\langle\Delta_{\pi^j}^\bullet\rangle). \tag{1}$$

As in the proof of Theorem 2.4, the first two and hence the third pro-simplicial spectrum in this fiber sequence are uniformly bounded below by $-\dim(A_0)$. Passing to geometric realizations and then to the limit “ \lim_j ”, the first term gets identified with $K^{\pi^\infty}(A_0 \text{ on } (\pi))$. For the second term, we compute

$$\begin{aligned} \text{“}\lim_j\text{” } |K^{\text{cont}}(A_0\langle\Delta_{\pi^j}^\bullet\rangle)| &= \text{“}\lim_j\text{” } | \text{“}\lim_n\text{” } K(A_0\langle\Delta_{\pi^j}^\bullet\rangle/(\pi^n)) | \\ &\simeq \text{“}\lim_j\text{” } \text{“}\lim_n\text{” } |K(A_0\langle\Delta_{\pi^j}^\bullet\rangle/(\pi^n))| && \text{(Lemma 2.8)} \\ &\simeq \text{“}\lim_n\text{” } \text{“}\lim_j\text{” } |K(A_0\langle\Delta_{\pi^j}^\bullet\rangle/(\pi^n))| \\ &\simeq \text{“}\lim_n\text{” } K(A_0/(\pi^n)) && (*) \\ &= K^{\text{cont}}(A_0). \end{aligned}$$

Here, the equivalence (*) follows from the isomorphism of pro-simplicial rings “ \lim_j ” $A_0\langle\Delta_{\pi^j}^\bullet\rangle/(\pi^n) \cong A_0/(\pi^n)$, where we view the right-hand term as a constant pro-simplicial ring. Thus, the weak fiber sequence (1) gives rise to the weak fiber sequence

$$K^{\pi^\infty}(A_0 \text{ on } (\pi)) \rightarrow K^{\text{cont}}(A_0) \rightarrow \text{“}\lim_j\text{” } K^{\text{cont}}(A\langle\Delta_{\pi^j}^\bullet\rangle).$$

Comparing with the weak fiber sequence of Theorem 2.4, we obtain the requested weak equivalence. \square

There is a more general version of the fundamental fiber sequence involving a Raynaud model X of the Tate ring $A = A_0[\frac{1}{\pi}]$. We use this to prove descent for analytic K -theory in §4. Let $p: X \rightarrow \text{Spec}(A_0)$ be an admissible morphism, that is, a proper morphism of schemes which is an isomorphism over $\text{Spec}(A)$. Let

$$K^{\text{cont}}(X) = \text{“}\lim_j\text{” } K(X \otimes_{A_0} A_0/(\pi^j)).$$

THEOREM 2.10. *Let A_0 be a complete π -adic ring which is Noetherian and of finite Krull dimension, and let $A = A_0[\frac{1}{\pi}]$. Let $p: X \rightarrow \text{Spec}(A_0)$ be an admissible morphism. Then there is a weak fiber sequence of pro-spectra*

$$K^{\pi^\infty}(X \text{ on } (\pi)) \rightarrow K^{\text{cont}}(X) \rightarrow K^{\text{an}}(A).$$

Proof. Thanks to Theorem 2.4, it suffices to show that in $\text{Pro}(\mathbf{Sp})$ there is a weakly Cartesian square

$$\begin{array}{ccc} K^{\pi^\infty}(A_0 \text{ on } (\pi)) & \longrightarrow & K^{\text{cont}}(A_0) \\ \downarrow & & \downarrow \\ K^{\pi^\infty}(X \text{ on } (\pi)) & \longrightarrow & K^{\text{cont}}(X). \end{array} \tag{2}$$

We have a commutative diagram

$$\begin{array}{ccccc}
 K^{\pi^\infty}(A_0 \text{ on } (\pi)) & \longrightarrow & K^{\pi^\infty}(A_0) & \longrightarrow & KH(A) \\
 \downarrow & & \downarrow & & \downarrow \cong \\
 K^{\pi^\infty}(X \text{ on } (\pi)) & \longrightarrow & K^{\pi^\infty}(X) & \longrightarrow & KH(X[\frac{1}{\pi}])
 \end{array} \tag{3}$$

where the rows are fiber sequences in $\text{Pro}(\mathbf{Sp})$ provided by Lemma 2.3. Since the right vertical map is an equivalence, the left-hand square is Cartesian. Write $X/(\pi^m) = X \otimes_{A_0} A_0/(\pi^m)$. We claim that the square

$$\begin{array}{ccc}
 K^{\pi^\infty}(A_0) & \longrightarrow & \text{“lim”}_{j,m} K(A_0/(\pi^m)[\Delta_{\pi^j}]) \\
 \downarrow & & \downarrow \\
 K^{\pi^\infty}(X) & \longrightarrow & \text{“lim”}_{j,m} K(X/(\pi^m)[\Delta_{\pi^j}])
 \end{array} \tag{4}$$

in $\text{Pro}(\mathbf{Sp})$ is weakly Cartesian. For this, let $Q(p, j, m)$ be the square of spectra

$$\begin{array}{ccc}
 K(A_0[\Delta_{\pi^j}^p]) & \longrightarrow & K(A_0/(\pi^m)[\Delta_{\pi^j}^p]) \\
 \downarrow & & \downarrow \\
 K(X[\Delta_{\pi^j}^p]) & \longrightarrow & K(X/(\pi^m)[\Delta_{\pi^j}^p])
 \end{array}$$

and denote by $F(p, j, m)$ its total fiber. So the square (4) is

$$\text{“lim”}_{j,m} \text{colim}_{p \in \Delta^{\circ p}} Q(p, j, m).$$

By *pro cdh descent* [8, Th. A], the square of pro-spectra “lim” $_m Q(p, j, m)$ is weakly Cartesian for every p and every j , that is, “lim” $_m F(p, j, m) \simeq 0$. Note also that by Weibel’s K -dimension conjecture [8, Th. B], all spectra in $Q(p, j, m)$ are $(-d)$ -connective for $d = \dim(A_0)$, and hence $F(p, j, m)$ is $(-d - 2)$ -connective. The spectral sequence computing the homotopy groups of the geometric realization of a simplicial spectrum now implies that the pro-spectrum

$$\text{“lim”}_m \text{colim}_{p \in \Delta^{\circ p}} F(p, j, m)$$

is weakly contractible for every j . Taking “lim” $_j$ implies that (4) is weakly Cartesian.

Finally, consider the commutative square in $\text{Pro}(\mathbf{Sp})$:

$$\begin{array}{ccc}
 K(A_0/(\pi^m)) & \xrightarrow{\simeq} & \text{“lim”}_j K(A_0/(\pi^m)[\Delta_{\pi^j}]) \\
 \downarrow & & \downarrow \\
 K(X/(\pi^m)) & \xrightarrow{\simeq} & \text{“lim”}_j K(X/(\pi^m)[\Delta_{\pi^j}]).
 \end{array} \tag{5}$$

The horizontal maps in the above diagram are equivalences by the fact that the map $A_0/(\pi^m) \rightarrow \text{“lim”}_j A_0/(\pi^m)[\Delta_{\pi^j}]$ is an isomorphism of pro-simplicial rings and a similar fact with A_0 replaced by X .

Composing the left-hand square in (3), (4), and the horizontal inverse of (5), we obtain the desired weakly Cartesian square (2). \square

§3. Excision for analytic K-theory

The goal of this section is to establish excision for Milnor squares for analytic K-theory for Tate rings that admit a finite-dimensional, Noetherian ring of definition.

Recall that a *Milnor square* is a Cartesian square of rings of the form

$$\begin{CD} A @>>> \bar{A} \\ @V\phi VV @VVV \\ B @>>> \bar{B} \end{CD} \tag{6}$$

in which the horizontal maps are surjective. By a *Milnor square of complete Tate rings*, we mean a Milnor square in which all rings are complete Tate rings and all homomorphisms are continuous.

LEMMA 3.1. *Assume that (6) is a Milnor square of complete Tate rings. Then there exist rings of definition A_0, B_0, \bar{A}_0 , and \bar{B}_0 such that (6) restricts to a Milnor square*

$$\begin{CD} A_0 @>>> \bar{A}_0 \\ @V\phi VV @VVV \\ B_0 @>>> \bar{B}_0. \end{CD}$$

If A and B admit finite-dimensional Noetherian rings of definition, one can choose these rings of definition to be Noetherian and finite-dimensional as well.

Proof. First note that if A admits the Noetherian ring of definition A_0 and A_1 is another ring of definition containing A_0 , then A_1 is finite over A_0 and in particular Noetherian with $\dim(A_0) = \dim(A_1)$. Indeed, if $\pi \in A_0$ is a pseudo-uniformizer, then by boundedness A_1 is contained in the free A_0 -module $\pi^{-n}A_0$ for n big enough. Keeping this in mind, the second claim will be clear from the following construction.

Choose rings of definition A_0 and B_0 of A and B , respectively. Replacing B_0 by $B_0 \cdot \phi(A_0)$ if necessary, we may assume that ϕ restricts to a map $A_0 \rightarrow B_0$. Now, let \bar{A}_0 be the image of A_0 in \bar{A} , and let \bar{B}_0 be the image of B_0 in \bar{B} . Define $A_1 \subseteq A$ as the pullback

$$A_1 = B_0 \times_{\bar{B}_0} \bar{A}_0.$$

By Banach’s open mapping theorem (see [2] for a version for Tate rings), the continuous isomorphism $A \rightarrow B \times_{\bar{B}} \bar{A}$ is a homeomorphism. As $B_0 \subseteq B$ and $\bar{A}_0 \subseteq \bar{A}$ are bounded, this implies that $A_1 \subseteq A$ is bounded. As it also contains A_0 , it is open and hence a ring of definition. Replacing A_0 by A_1 , we have constructed the desired rings of definition. \square

THEOREM 3.2. *Let (6) be a Milnor square of complete Tate rings. Assume that A and B admit Noetherian rings of definition that are finite-dimensional. Then the induced square of pro-spectra*

$$\begin{array}{ccc}
 K^{\text{an}}(A) & \longrightarrow & K^{\text{an}}(\overline{A}) \\
 \downarrow & & \downarrow \\
 K^{\text{an}}(B) & \longrightarrow & K^{\text{an}}(\overline{B})
 \end{array} \tag{an}$$

is weakly Cartesian.

Proof. Using nil-invariance of K^{an} , we will reduce this to pro-excision for K^{cont} and K^{π^∞} of adic rings, using Theorem 2.4.

Choose Noetherian, finite-dimensional rings of definition as in Lemma 3.1, and let $I_0 = \ker(A_0 \rightarrow \overline{A}_0)$, $J_0 = \ker(B_0 \rightarrow \overline{B}_0)$, so that $I_0 \cong J_0$ via ϕ . For every $m \geq 1$,

$$(A_0/I_0^m)[\frac{1}{\pi}] \rightarrow (A_0/I_0)[\frac{1}{\pi}] \cong \overline{A}$$

is then a nilpotent extension of Tate rings, and hence

$$K^{\text{an}}((A_0/I_0^m)[\frac{1}{\pi}]) \rightarrow K^{\text{an}}(\overline{A})$$

is a weak equivalence by [7, Prop. 6.17]. From Theorem 2.4, we therefore deduce a weak fiber sequence

$$\text{“}\lim_m\text{” } K^{\pi^\infty}(A_0/I_0^m \text{ on } (\pi)) \rightarrow \text{“}\lim_m\text{” } K^{\text{cont}}(A_0/I_0^m) \rightarrow K^{\text{an}}(\overline{A})$$

and similarly for B_0, J_0 . So we get a weak fiber sequence of squares of pro-spectra

$$\boxed{\pi^\infty} \rightarrow \boxed{\text{cont}} \rightarrow \boxed{\text{an}}.$$

The theorem thus follows from the following two pro-excision Claims 3.3 and 3.4. □

CLAIM 3.3. *The square of pro-spectra*

$$\begin{array}{ccc}
 K^{\text{cont}}(A_0) & \longrightarrow & \text{“}\lim_m\text{” } K^{\text{cont}}(A_0/I_0^m) \\
 \downarrow \phi & & \downarrow \\
 K^{\text{cont}}(B_0) & \longrightarrow & \text{“}\lim_m\text{” } K^{\text{cont}}(B_0/J_0^m)
 \end{array} \tag{\text{cont}}$$

is weakly Cartesian.

CLAIM 3.4. *The square of pro-spectra*

$$\begin{array}{ccc}
 K^{\pi^\infty}(A_0 \text{ on } (\pi)) & \longrightarrow & \text{“}\lim_m\text{” } K^{\pi^\infty}(A_0/I_0^m \text{ on } (\pi)) \\
 \downarrow \phi & & \downarrow \\
 K^{\pi^\infty}(B_0 \text{ on } (\pi)) & \longrightarrow & \text{“}\lim_m\text{” } K^{\pi^\infty}(B_0/J_0^m \text{ on } (\pi))
 \end{array} \tag{\pi^\infty}$$

is weakly Cartesian.

Proof of Claim 3.3. As $I_0 \xrightarrow{\sim} J_0$ via ϕ , also $I_0^m \xrightarrow{\sim} J_0^m$. As π is a nonzero divisor on A_0/I_0 , it is also a nonzero divisor on A_0/I_0^m for all m , and similarly on B_0/J_0^m . It follows

from these two observations that for every pair of positive integers n, m , we have a Milnor square

$$\begin{CD} A_0/(\pi^n) @>>> A_0/(\pi^n, I_0^m) \\ @V\phi VV @VVV \\ B_0/(\pi^n) @>>> B_0/(\pi^n, J_0^m). \end{CD} \tag{7}$$

Pro-excision for Noetherian rings (see [13, Cor. 0.4]; see also [9, §2.4] for an account) implies that the square of K -theory pro-spectra

$$\text{“}\varinjlim\text{”}_m K((7))$$

is weakly contractible for every n . Taking the limit over n gives the claim. □

Proof of Claim 3.4. From pro-excision for Noetherian rings, applied to the Milnor squares

$$\begin{CD} A_0[\Delta_{\pi^j}^p] @>>> A_0/I_0^m[\Delta_{\pi^j}^p] \\ @V\phi VV @VVV \\ B_0[\Delta_{\pi^j}^p] @>>> B_0/J_0^m[\Delta_{\pi^j}^p], \end{CD} \tag{8}$$

we deduce that for fixed p and j , the square of pro-spectra

$$\text{“}\varinjlim\text{”}_m K((8) \text{ on } (\pi))$$

is weakly Cartesian. As in the proof of Theorem 2.10, using finite dimensionality and Weibel’s conjecture, we deduce that we can pass to geometric realizations, and deduce the claim. □

§4. Descent and applications

4.1 Descent for analytic K -theory

Let (A, A^+) be an affinoid ring in the sense of [4, §3], and let $X = \text{Spa}(A, A^+)$. We assume that A is Tate and that it admits a Noetherian, finite-dimensional ring of definition A_0 . If $U \subseteq X$ is a rational subdomain, then the same conditions hold for the affinoid ring $(\mathcal{O}_X(U), \mathcal{O}_X(U)^+)$. We fix a pseudo-uniformizer π of A . Then (the image of) π is also a pseudo-uniformizer for every Tate ring $\mathcal{O}_X(U)$. For ease of notation, we set

$$K^{\text{an}}(U) := K^{\text{an}}(\mathcal{O}_X(U)).$$

Let $f_1, \dots, f_n \in A$ be elements generating A as an ideal. These give rise to the standard rational covering $\mathcal{U} = (U_i)_{1 \leq i \leq n}$ of X , where

$$U_i = X \left(\frac{f_1, \dots, f_n}{f_i} \right).$$

Write

$$K^{\text{an}}(|\mathcal{U}_\bullet|) := \varinjlim_{p \in \Delta} K^{\text{an}}(\mathcal{U}_p), \tag{9}$$

where \mathcal{U}_\bullet is the Čech nerve of the covering \mathcal{U} .

THEOREM 4.1. *In the above situation, the natural map*

$$K^{\text{an}}(A) \rightarrow K^{\text{an}}(|\mathcal{U}_\bullet|)$$

is a weak equivalence of pro-spectra.

Proof. Fix a finite-dimensional, Noetherian ring of definition A_0 of A . Multiplying by a power of π , we may assume that the f_i belong to A_0 . Let $I \subseteq A_0$ be the ideal generated by f_1, \dots, f_n , and let

$$Y = \text{Proj}\left(\bigoplus_{k \geq 0} I^k\right)$$

be the blowup of $\text{Spec}(A_0)$ along the ideal I . Then $Y \rightarrow \text{Spec}(A_0)$ is an admissible morphism. Let $V_i = D_+(f_i) \subseteq Y$. Then $\mathcal{V} = (V_i)_{1 \leq i \leq n}$ is an affine open covering of Y . The π -adic completion $\mathcal{O}_Y(V_i)^\wedge$ can be canonically identified with a ring of definition of the Tate ring $\mathcal{O}_X(U_i)$, and similarly for the higher intersections $V_{i_0} \cap \dots \cap V_{i_p}$.

Consider the following diagram of pro-spectra

$$\begin{CD} K^{\pi^\infty}(Y \text{ on } (\pi)) @>>> K^{\text{cont}}(Y) @>>> K^{\text{an}}(A) \\ @VVV @VVV @VVV \\ K^{\pi^\infty}(|\mathcal{V}_\bullet| \text{ on } (\pi)) @>>> K^{\text{cont}}(|\mathcal{V}_\bullet|) @>>> K^{\text{an}}(|\mathcal{U}_\bullet|), \end{CD} \tag{10}$$

where $K^{\text{cont}}(|\mathcal{V}_\bullet|)$ and $K^{\pi^\infty}(|\mathcal{V}_\bullet| \text{ on } (\pi))$ are defined as in (9). The top line is a weak fiber sequence by Theorem 2.10. Note that the coverings \mathcal{U} and \mathcal{V} consist of n open subsets of X and Y , respectively, and thus their Čech nerves are n -skeletal. This implies that the limit

$$K^{\text{cont}}(|\mathcal{V}_\bullet|) = \lim_{[p] \in \Delta} K^{\text{cont}}(\mathcal{V}_p)$$

in $\text{Pro}(\mathbf{Sp})$ can be computed level-wise (see, e.g., [7, Lem. 2.1]) and similarly for $K^{\pi^\infty}(|\mathcal{V}_\bullet| \text{ on } (\pi))$. From Zariski descent for nonconnective K -theory (see [14, Th. 8.1]), we thus deduce that the left two vertical maps in (10) are equivalences of pro-spectra.

Let

$$V'_i = \text{Spec}(\mathcal{O}_Y(V_i)^\wedge)$$

and similarly for the higher intersections. By Thomason’s excision [14, Prop. 3.19], we have an equivalence

$$K^{\pi^\infty}(|\mathcal{V}_\bullet| \text{ on } (\pi)) \simeq K^{\pi^\infty}(|\mathcal{V}'_\bullet| \text{ on } (\pi)),$$

and we clearly also have $K^{\text{cont}}(|\mathcal{V}_\bullet|) \simeq K^{\text{cont}}(|\mathcal{V}'_\bullet|)$. Hence, we deduce from Theorem 2.4 that the lower horizontal line in (10) is also a fiber sequence. We conclude that the right vertical map is a weak equivalence, as was to be shown. \square

4.2 Globalization

In order to globalize the construction of analytic K -theory, we first recall some sheaf theory. Let C be a small category. We denote by

$$\mathbf{PSh}(C) = \text{Fun}(C^{\text{op}}, \mathbf{Spc})$$

the ∞ -category of presheaves of spaces on C and by $y: C \rightarrow \mathbf{PSh}(C)$ the Yoneda embedding. If F and U are presheaves, we write

$$F(U) = \text{Map}_{\mathbf{PSh}(C)}(U, F).$$

Note that by the Yoneda lemma, we have $F(X) = F(y(X))$ for every object X of C .

If \mathbf{D} is any complete ∞ -category, then composition with the Yoneda embedding induces an equivalence

$$\text{Fun}^{lim}(\mathbf{PSh}(C)^{op}, \mathbf{D}) \xrightarrow{\simeq} \text{Fun}(C^{op}, \mathbf{D}), \tag{11}$$

where $\text{Fun}^{lim}(\mathbf{PSh}(C)^{op}, \mathbf{D}) \subseteq \text{Fun}(\mathbf{PSh}(C)^{op}, \mathbf{D})$ denotes the full subcategory spanned by the limit preserving functors (see [10, Th. 5.1.5.6]).

We now assume that C is equipped with a (Grothendieck) topology. There is a bijection between sieves on an object $X \in C$ and subobjects $U \hookrightarrow y(X)$ of $y(X)$ in $\mathbf{PSh}(C)$ (see [10, Prop. 6.2.2.5]). For simplicity, we call the latter sieves, too. By definition, a presheaf $F \in \mathbf{PSh}(C)$ is a sheaf if and only if the natural map

$$F(X) \rightarrow F(U)$$

is an equivalence for every $X \in C$ and every covering sieve $U \hookrightarrow y(X)$ of X . We denote by

$$\mathbf{Sh}(C) \subseteq \mathbf{PSh}(C)$$

the full subcategory consisting of sheaves. The inclusion $\mathbf{Sh}(C) \hookrightarrow \mathbf{PSh}(C)$ admits a left adjoint $L: \mathbf{PSh}(C) \rightarrow \mathbf{Sh}(C)$ called *sheafification* (see [10, Prop. 5.5.4.15 and Lem. 6.2.2.7]). We say that a functor $F \in \text{Fun}(C^{op}, \mathbf{D})$ is a (\mathbf{D} -valued) sheaf if $F(X) \rightarrow F(U)$ is an equivalence in \mathbf{D} for every X and covering sieve $U \hookrightarrow y(X)$ as above. Here, $F(U)$ is defined using (11) above. By [10, Prop. 5.5.4.20], composition with L induces a fully faithful functor

$$\text{Fun}^{lim}(\mathbf{Sh}(C)^{op}, \mathbf{D}) \rightarrow \text{Fun}^{lim}(\mathbf{PSh}(C)^{op}, \mathbf{D}) \simeq \text{Fun}(C^{op}, \mathbf{D}),$$

whose essential image consists precisely of the \mathbf{D} -valued sheaves.

We now return to analytic K -theory of Tate rings. Fix a complete affinoid Tate ring (R, R^+) such that R admits a finite-dimensional, Noetherian ring of definition R_0 , and let $S = \text{Spa}(R, R^+)$. We denote by

$$\mathbf{Ad}_S^{\text{aff,ft}} \subseteq \mathbf{Ad}_S^{\text{sep,lft}} \subseteq \mathbf{Ad}_S^{\text{lft}},$$

the categories of affinoid adic spaces of finite type over S , of adic spaces separated and locally of finite type over S , and of adic spaces locally of finite type over S , respectively (see [5, §3] and [6, §1.2 and 1.3] for these notions). Note that for any $X \in \mathbf{Ad}_S^{\text{lft}}$ and any open affinoid $U = \text{Spa}(A, A^+) \subseteq X$, A admits a finite-dimensional, Noetherian ring of definition. Indeed, by [5, Prop. 3.6], the map $(R, R^+) \rightarrow (A, A^+)$ is topologically of finite type. By Lemma 3.3(iii) there, $R \rightarrow A$ factors through a continuous open surjection $R\langle x_1, \dots, x_n \rangle \rightarrow A$, and we can take the image of $R_0\langle x_1, \dots, x_n \rangle$ in A as a ring of definition.

We equip the categories of adic spaces above with the usual topologies (see [4, §2]). We then have the following lemma.

LEMMA 4.2. *The inclusion $\mathbf{Ad}_S^{\text{aff,ft}} \subseteq \mathbf{Ad}_S^{\text{lft}}$ induces an equivalence*

$$\mathbf{Sh}(\mathbf{Ad}_S^{\text{lft}}) \simeq \mathbf{Sh}(\mathbf{Ad}_S^{\text{aff,ft}}).$$

Proof. This follows by applying [3, Lem. C.3] to both inclusions $\mathbf{Ad}_S^{\text{aff,ft}} \subseteq \mathbf{Ad}_S^{\text{sep,ft}} \subseteq \mathbf{Ad}_S^{\text{ft}}$. Here, we use that every $X \in \mathbf{Ad}_S^{\text{ft}}$ has a covering $\{U_i \rightarrow X\}$ such that all intersections $U_{i_0} \cap \dots \cap U_{i_n}$ are separated, and any separated X has such a covering with all intersections affinoid. \square

COROLLARY 4.3. *The presheaf*

$$K^{\text{an}} : \mathbf{Ad}_S^{\text{aff,ft,op}} \rightarrow \text{Pro}(\mathbf{Sp}^+), \quad \text{Spa}(A, A^+) \mapsto K^{\text{an}}(A),$$

is a sheaf. In particular, it extends essentially uniquely to a sheaf with values in $\text{Pro}(\mathbf{Sp}^+)$ on the category $\mathbf{Ad}_S^{\text{ft}}$.

In particular, we can now make sense of the analytic K -theory $K^{\text{an}}(X)$ for every adic space X locally of finite type over the base S .

Proof. In view of the general discussion at the beginning of this subsection, the second assertion follows from the first one and Lemma 4.2. Now, we prove the first one. For each affinoid X , we define a family of sieves $\tau(X)$ on X as follows: a sieve $U \hookrightarrow y(X)$ belongs to $\tau(X)$ if and only if the natural map

$$K^{\text{an}}(X') \rightarrow K^{\text{an}}(U \times_{y(X)} y(X'))$$

is an equivalence in $\text{Pro}(\mathbf{Sp}^+)$ for every morphism $X' \rightarrow X$ in $\mathbf{Ad}_S^{\text{aff,ft}}$. Then the association $X \rightsquigarrow \tau(X)$ defines a topology on $\mathbf{Ad}_S^{\text{aff,ft}}$ (see [3, Prop. C.1]). Since the topology on $\mathbf{Ad}_S^{\text{aff,ft}}$ is generated by rational coverings, it suffices to show that the sieve generated by a rational covering of X belongs to $\tau(X)$. Since rational coverings are stable under pullback, it suffices to show that

$$K^{\text{an}}(X) \rightarrow K^{\text{an}}(|U_\bullet|)$$

is an equivalence, where $|U_\bullet| = \text{colim}_{[p] \in \Delta^{\text{op}}} y(U_{[p]})$ is the realization of the Čech nerve of a rational covering of X . Note here that $|U_\bullet| \hookrightarrow y(X)$ is precisely the sieve generated by that covering. This equivalence is a consequence of Theorem 4.1. \square

4.3 Properties of analytic K -theory

In this subsection, we record some immediate consequences. We keep the assumptions on the affinoid adic base space S as in the previous subsection.

PROPOSITION 4.4. (\mathbb{A}^1 -invariance) *Let X be an adic space locally of finite type over S , and let \mathbb{A}_X^1 denote the adic affine line over X . Then the natural map*

$$K^{\text{an}}(X) \rightarrow K^{\text{an}}(\mathbb{A}_X^1)$$

is an equivalence in $\text{Pro}(\mathbf{Sp}^+)$.

Proof. It suffices to check this when $X = \text{Spa}(A, A^+)$ is affinoid. Let $X\langle t \rangle = \text{Spa}(A\langle t \rangle, A^+\langle t \rangle)$. The sheaf (of spaces) $y(\mathbb{A}_X^1)$ is equivalent to $\text{colim}_{t \rightarrow \pi t} y(X\langle t \rangle)$. Hence,

$$K^{\text{an}}(\mathbb{A}_X^1) \simeq \lim_{t \rightarrow \pi t} K^{\text{an}}(X\langle t \rangle) \simeq K^{\text{an}}(X),$$

where the second equivalence follows from Corollary 2.7. \square

Recall that the K -module structure of K^{an} as a functor on affinoids provides a natural transformation $\smile t: K^{\text{an}} \rightarrow \Omega K^{\text{an}}(_ \langle t, t^{-1} \rangle)$. It induces a natural transformation on the extension of K^{an} to all adic spaces in $\mathbf{Ad}_S^{\text{ft}}$.

PROPOSITION 4.5. (*Projective line*) Let X be an adic space locally of finite type over S , and denote by \mathbb{P}_X^1 the adic projective line over X . We have a natural weak equivalence

$$K^{\text{an}}(\mathbb{P}_X^1) \simeq K^{\text{an}}(X) \oplus K^{\text{an}}(X).$$

Proof. We may again assume that X is affinoid. Consider the covering of \mathbb{P}_X^1 given by $X\langle t \rangle, X\langle t^{-1} \rangle$. Denote the projection $\mathbb{P}_X^1 \rightarrow X$ by p and let $q := p|_{X\langle t, t^{-1} \rangle}$. Since K^{an} is a sheaf, the upper line in the following diagram is a fiber sequence.

$$\begin{array}{ccccc} \Omega K^{\text{an}}(X\langle t, t^{-1} \rangle) & \longrightarrow & K^{\text{an}}(\mathbb{P}_X^1) & \longrightarrow & K^{\text{an}}(X\langle t \rangle) \oplus K^{\text{an}}(X\langle t^{-1} \rangle) \\ q^* \uparrow & & p^* \uparrow & & \simeq \uparrow \\ \Omega \Gamma K^{\text{an}}(X) & \longrightarrow & K^{\text{an}}(X) & \longrightarrow & K^{\text{an}}(X\langle t \rangle) \oplus K^{\text{an}}(X\langle t^{-1} \rangle) \end{array}$$

Here,

$$\Gamma K^{\text{an}}(X) = K^{\text{an}}(X\langle t \rangle) \sqcup_{K^{\text{an}}(X)} K^{\text{an}}(X\langle t^{-1} \rangle).$$

The lower line is a fiber sequence by definition of $\Gamma K^{\text{an}}(X)$. Since p^* is split, we have an equivalence $K^{\text{an}}(\mathbb{P}_X^1) \simeq K^{\text{an}}(X) \oplus \text{cofib}(p^*)$. From the diagram, we get a weak equivalence $\text{cofib}(q^*) \simeq \text{cofib}(p^*)$. By definition of ΛK^{an} (see §2.2), we have an equivalence $\text{cofib}(q^*) \simeq \Lambda K^{\text{an}}(A)$. By Corollary 2.6, the composite

$$K^{\text{an}}(X) \xrightarrow{\simeq} \Omega K^{\text{an}}(X\langle t, t^{-1} \rangle) \rightarrow \Lambda K^{\text{an}}(X) \simeq \text{cofib}(q^*) \simeq \text{cofib}(p^*)$$

is a weak equivalence. Thus, this map together with p^* induces the desired weak equivalence $K^{\text{an}}(X) \oplus K^{\text{an}}(X) \xrightarrow{\simeq} K^{\text{an}}(\mathbb{P}_X^1)$. □

We can globalize Theorem 2.10 as follows. Recall that (R, R^+) denotes an affinoid Tate ring such that R admits a Noetherian, finite-dimensional ring of definition R_0 , $\pi \in R_0$ is a pseudo-uniformizer, and $S = \text{Spa}(R, R^+)$. Denote by $\mathbf{Sch}_{R_0}^{\text{ft}}$ the category of R_0 -schemes of finite type. There is a functor

$$\mathbf{Sch}_{R_0}^{\text{ft}} \rightarrow \mathbf{Ad}_S^{\text{ft}}, \quad X \mapsto X^{\text{ad}},$$

which on affine schemes is given by $\text{Spec}(A_0) \mapsto \text{Spa}(A, A^\circ)$ with $A = A_0^\wedge[\frac{1}{\pi}]$ and $A^\circ \subseteq A$ the subring of power-bounded elements (compose the π -adic completion functor, the functor t from [5, Prop. 4.1], which sends $\text{Spf}(A_0^\wedge)$ to $\text{Spa}(A_0^\wedge, A_0^\wedge)$, and the functor $Y \mapsto Y(\pi \neq 0)$).

THEOREM 4.6. For X a scheme of finite type over R_0 , there is a weak fiber sequence of pro-spectra

$$K^{\pi^\infty}(X \text{ on } (\pi)) \rightarrow K^{\text{cont}}(X) \rightarrow K^{\text{an}}(X^{\text{ad}}).$$

If there exists an admissible morphism $\tilde{X} \rightarrow X$ with \tilde{X} regular, then there is a weak fiber sequence

$$K(X \text{ on } (\pi)) \rightarrow K^{\text{cont}}(X) \rightarrow K^{\text{an}}(X^{\text{ad}}).$$

Proof. We equip $\mathbf{Sch}_{R_0}^{\text{ft}}$ with the Zariski topology. Considered as functors on $\mathbf{Sch}_{R_0}^{\text{ft}, \text{op}}$, all pro-spectra appearing in the statement of the theorem are sheaves (see the proof of Theorem 4.1 for the cases of $K^{\pi^\infty}(_ \text{ on } (\pi))$ and K^{cont}). Since every (regular) $X \in \mathbf{Sch}_{R_0}^{\text{ft}}$

is a colimit of (regular) affine schemes in the category of sheaves $\mathbf{Sh}(\mathbf{Sch}_{R_0}^{\text{ft}})$, it suffices to check the first statement in the affine case. But this is Theorem 2.4.

If there exists an admissible morphism $\tilde{X} \rightarrow X$ with \tilde{X} regular, then [7, Cor. 4.7] implies that for every p , the map

$$K(X \text{ on } (\pi)) \rightarrow \varinjlim_j K(X \otimes_{R_0} R_0[\Delta_{\pi^j}^p] \text{ on } (\pi))$$

is a weak equivalence of pro-spectra. Using arguments similar to those in the proof of Theorem 2.4, Weibel's K -dimension conjecture implies that $K^{\pi^\infty}(X \text{ on } (\pi))$ is weakly equivalent to $K(X \text{ on } (\pi))$. This establishes the second claim. \square

REFERENCES

- [1] C. Dahlhausen, *Continuous K-theory and cohomology of rigid spaces*, preprint, [arXiv:1910.10437](https://arxiv.org/abs/1910.10437)
- [2] T. Henkel, *An open mapping theorem for rings which have a zero sequence of units*, preprint, [arXiv:1407.5647](https://arxiv.org/abs/1407.5647)
- [3] M. Hoyois, *A quadratic refinement of the Grothendieck–Lefschetz–Verdier trace formula*, *Algebr. Geom. Topol.* **14** (2014), 3603–3658.
- [4] R. Huber, *Continuous valuations*, *Math. Z.* **212** (1993), 455–477.
- [5] R. Huber, *A generalization of formal schemes and rigid analytic varieties*, *Math. Z.* **217** (1994), 513–551.
- [6] R. Huber, *Étale Cohomology of Rigid Analytic Varieties and Adic Spaces*, *Aspects of Math.* **E30**, Friedr. Vieweg & Sohn, Braunschweig, 1996.
- [7] M. Kerz, S. Saito, and G. Tamme, *K-theory of non-archimedean rings. I*, *Doc. Math.* **24** (2019), 1365–1411.
- [8] M. Kerz, F. Strunk, and G. Tamme, *Algebraic K-theory and descent for blow-ups*, *Invent. Math.* **211** (2018), 523–577.
- [9] M. Land and G. Tamme, *On the K-theory of pullbacks*, *Ann. of Math. (2)* **190** (2019), 877–930.
- [10] J. Lurie, *Higher Topos Theory*, *Ann. of Math. Stud.* **170**, Princeton University Press, Princeton, NJ, 2009.
- [11] J. Lurie, *Higher Algebra*, book preprint (version September 2017), 2017. <http://www.math.ias.edu/lurie/>.
- [12] M. Morrow, *A historical overview of pro cdh descent in algebraic K-theory and its relation to rigid analytic varieties*, preprint, [arXiv:1612.00418](https://arxiv.org/abs/1612.00418)
- [13] M. Morrow, *Pro unitality and pro excision in algebraic K-theory and cyclic homology*, *J. Reine Angew. Math.* **736** (2018), 95–139.
- [14] R. Thomason and T. Trobaugh, “Higher algebraic K-theory of schemes and of derived categories” in *The Grothendieck Festschrift, Volume III*, *Progr. Math.* **88**, Birkhäuser, Boston, MA, 1990, 247–435.

Moritz Kerz
Fakultät für Mathematik
Universität Regensburg
93040 Regensburg
Germany
moritz.kerz@ur.de

Shuji Saito
Graduate School of Mathematical Sciences
University of Tokyo
3-8-1 Komaba, Tokyo
Japan
sshuji@msb.biglobe.ne.jp

Georg Tamme
Institut für Mathematik, Fachbereich 08
Johannes Gutenberg-Universität Mainz
55099 Mainz
Germany
georg.tamme@uni-mainz.de