

# Reconstructing caribou seasonal biogeography in Little Ice Age (late Holocene) Western Alaska using intra-tooth strontium and oxygen isotope analysis

Ciara Gigueux<sup>a,\*</sup>, Vaughan Grimes<sup>b</sup>, Thomas Tütken<sup>c</sup>, Rick Knecht<sup>a</sup>, Kate Britton<sup>a,d</sup>

<sup>a</sup> Department of Archaeology, University of Aberdeen, St. Mary's Building, Elphinstone Road, Aberdeen AB24 3UF, UK

<sup>b</sup> Department of Archaeology, Memorial University, St. John's, NL A1C 5S7, Canada

<sup>c</sup> Institut für Geowissenschaften, AG für Angewandte und Analytische Paläontologie, Universität Mainz, J.-J. Becherweg 21, 55128 Mainz, Germany

<sup>d</sup> Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany

## ARTICLE INFO

### Keywords:

Isotopes  
Enamel  
Rangifer  
Migration  
Ecology  
Climate

## ABSTRACT

The palaeobiogeography of key prey-species can provide valuable insights into animal-human interactions, human subsistence activities and landscape use in the past. In many contemporary indigenous Arctic societies, caribou (*Rangifer tarandus* spp.) are an important seasonal subsistence species, and recent climatic shifts have influenced the seasonal and spatial distribution and migrations of herds. The impact of larger scale climatic change on this species, such as that experienced during the Little Ice Age (LIA), is not known, but may provide vital clues about future variability. Here we present sequential strontium ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) and oxygen ( $\delta^{18}\text{O}_{\text{CO}_3}$ ) isotope data from archaeological caribou tooth enamel from Nunalleq, a 15th to 17th century AD pre-contact Yup'ik village site in Western Alaska, to reconstruct caribou movement patterns in this region during the LIA. The results of these analyses highlight variation in ranging habits over the period of time that the site was occupied, and indicate different ranging behaviours in the region in the past compared to modern herds in the area today. The isotopic data presented here complement the wealth of data derived from other research at Nunalleq, illuminating the influence of changing climatic conditions on prey-species palaeoecology and human-animal interactions at the site.

## 1. Introduction

Isotopic analyses of sequentially sampled, incrementally-formed tissues are increasingly being utilised in archaeology and palaeoecology to identify and reconstruct seasonal dietary and migratory behaviours in wild animal species in the past (e.g. Britton et al., 2011; Feranec and MacFadden, 2000; Henton et al., 2017; Julien et al., 2012; Metcalfe and Longstaffe, 2014; Pellegrini et al., 2008; Price et al., 2017). Incrementally-developed tissues, such as tooth enamel and dentine, are formed over a relatively short period of time with no remodelling so the sequential sampling of these materials can provide time-series isotopic profiles. In this way, the episodic nature of isotopic variations measured within the tissues – brought about by variations in isotopic inputs related to changes in diet, ingested water, or location during tooth formation – can be established. Strontium isotope measurements in teeth, for example, are related to (geological) location during formation (Bentley, 2006), and – given that oxygen isotope ratios of precipitation

vary seasonally at mid-/high-latitudes – oxygen isotope data can be utilised alongside strontium to ‘anchor’ data within a seasonal context. The utility of intra-tooth strontium and oxygen isotope analysis for the reconstruction of seasonal migratory behaviour in caribou/reindeer (*Rangifer tarandus*) has been recently demonstrated using migratory modern caribou (Britton et al., 2009; Britton, 2010), and has since been applied to European Middle and Late Palaeolithic reindeer (Britton et al., 2011; Price et al., 2017).

When focused on archaeologically-important prey-species, isotope zooarchaeology (the isotope analysis of archaeofaunal remains) studies not only serve to reconstruct the seasonal movements of important extinct and ancestral extant taxa, but also offer insights into animal-human interactions, animal landscape use and past human subsistence strategies. Isotope zooarchaeology also has the potential to allow archaeologists and palaeoecologists to explore changes in animal ecological behaviours over time periods not achievable in modern ecological studies. The influence of modern-day climate and environmental

\* Corresponding author.

E-mail addresses: [c.gigueux@abdn.ac.uk](mailto:c.gigueux@abdn.ac.uk) (C. Gigueux), [vgrimes@mun.ca](mailto:vgrimes@mun.ca) (V. Grimes), [tuetken@uni-mainz.de](mailto:tuetken@uni-mainz.de) (T. Tütken), [r.knecht@abdn.ac.uk](mailto:r.knecht@abdn.ac.uk) (R. Knecht), [k.britton@abdn.ac.uk](mailto:k.britton@abdn.ac.uk) (K. Britton).

<https://doi.org/10.1016/j.jasrep.2017.10.043>

Received 13 June 2017; Received in revised form 15 September 2017; Accepted 28 October 2017

Available online 14 November 2017

2352-409X/ © 2017 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

fluctuations, including anthropogenically-induced ones, on prey-species ecology, and specifically their migrations, has been observed for a number of modern wild animal species including terrestrial and marine mammals and birds (e.g. Blois and Hadly, 2009; Post et al., 2009; Root et al., 2003; Tape et al., 2016). Considering this, and other factors, the ecological behaviours of archaeological species (even in extant species) should not simply be assumed using modern analogues (see Britton, in press). Furthermore, the isotope zooarchaeological record, particularly within the context of large scale climatic shifts, can inform on continuity and change in faunal palaeoecology and the impact of environmental changes on faunal species, and illuminate species adaptability and niche plasticity or conservatism (see Britton, in press). To date, however, few studies have utilised these approaches to assess behavioural changes, though a limited number of studies have sought to demonstrate changes in palaeoecological behaviours over time at a single site- or on a regional/population-specific level (e.g. Britton, in press; Price et al., 2017). Further studies are warranted to better understand the impact of environmental change on the behaviour of medium- and large-sized herbivores, both in the past and in the future, particularly given the rate and scale of present-day global warming-induced environmental instability.

Caribou and reindeer (*Rangifer tarandus*) are amongst the most important seasonal subsistence taxa in the circumpolar North, particularly for contemporary Indigenous communities. North American caribou have both sedentary and migratory ecotypes; sedentary caribou ‘space out’, moving only during calving to avoid predation, whilst migratory caribou ‘space away’, undertaking long-range seasonal movements usually on a North-South trajectory (Bergerud, 1988:77). Migratory herds demonstrate cyclical movements, showing fidelity to calving sites and geographic ranges (Gunn and Miller, 1986; Skoog, 1968), although the size and actual location of the range can vary depending on herd numbers, forage abundance, predation and modern infrastructure (Dahle et al., 2008; Gunn and Miller, 1986; Hemming, 1971; Hinzman et al., 2005). Modern-day climatic shifts have also been shown to influence caribou seasonal movements and range use (e.g. Abraham et al., 2012; Brotton and Wall, 1997; Jacobs et al., 1996; Masood et al., 2017; Weladji et al., 2002). Extrinsic markers such as telemetry, satellite transmitters and aerial photography can provide detailed information on caribou herd range use and movements (Hobson, 1999), however other methods – such as isotope analysis – are required to reconstruct caribou migratory habits and range use in the recent historical and archaeological past.

Here, we present coupled  $^{87}\text{Sr}/^{86}\text{Sr}$  and  $\delta^{18}\text{O}_{\text{CO}_3}$  isotopic measurements of sequentially-sampled *Rangifer* tooth enamel from the late Holocene site of Nunalleq in the Yukon-Kuskokwim Delta of Western Alaska. The pre-contact Yup'ik village site was primarily occupied between the 15th to 17th centuries (Britton et al., 2016; Ledger et al., 2016), during a period of global climatic disruption known as the Little Ice Age. The research goals of our study are: (i) to identify if these caribou undertook seasonal migrations in the past; (ii) to identify any differences in calving ground area and/or range use in the past compared to the region's present-day caribou population; and (iii) to utilise this new isotopic evidence for the (possible) seasonal movements of caribou in the Yukon-Kuskokwim Delta to better understand the nature of caribou exploitation at the site.

## 2. Reconstructing faunal migration using stable isotope analysis

Initial applications of stable isotope analysis demonstrated that the isotopic ratios of foods consumed during life are reflected in the isotopic ratios of body tissues (DeNiro and Epstein, 1978, 1981). Subsequent studies introduced and furthered the notion of applying the technique to reconstruct human palaeodiet (e.g. Tauber, 1981; Chisholm et al., 1982; Schoeninger et al., 1983). Since these initial studies, the use of isotope analysis in archaeology has expanded substantially, with studies now also encompassing the reconstruction of

past animal diets (e.g. Balasse et al., 2006; Craig et al., 2006; Feranec et al., 2009; Sayle et al., 2016); human and animal mobility (e.g. Bentley and Knipper, 2005; Hoppe et al., 1999; Montgomery et al., 2007; Price et al., 2000); human and animal behaviours such as weaning (e.g. Britton et al., 2015; Fuller et al., 2003) and birth seasonality (e.g. Balasse et al., 2003; Buchan et al., 2016), and the reconstruction of palaeoenvironments and palaeoclimate (e.g. Domingo et al., 2015; Fricke et al., 1998; Hedges et al., 2004; Stevens et al., 2014; Tütken et al., 2008; Wißing et al., 2015). Early studies primarily utilised ‘bulk’ measurements of bones and teeth, giving averaged values of different ontogenetic periods (depending on the formation and/or remodelling time of the hard tissue in question). The desire for higher-resolution isotopic data, however, has led to an increase in the sequential sampling of incrementally-formed tissues such as enamel, dentine, and antler.

The accumulative nature of enamel growth, whereby enamel is laid down in successive layers and then mineralises, occurs over a relatively conservative period during sub-adulthood. Unlike bone, these incremental layers of tissue do not remodel over time, such that the isotopic composition of the diet consumed during their growth is fixed within these tissues. Intra-tooth isotopic analysis often involves sampling an enamel face horizontally across these bands of growth, therefore providing time-series isotopic data. Some biases, such as isotopic attenuation due to sample homogenisation across multiple phases of growth and within-tooth variations in growth-rates, can occur during sampling (Balasse, 2003; Bendrey et al., 2015; Zazzo et al., 2006, 2012). In archaeofaunal studies, such sampling has enabled a deeper understanding of seasonal dietary behaviour (e.g. Balasse et al., 2006; Feranec et al., 2009), animal husbandry practices (e.g. Balasse et al., 2012; Balasse and Tresset, 2002) and seasonal mobility (e.g. Britton et al., 2011; Julien et al., 2012; Widga et al., 2010). The period of time represented depends on the rate of tooth growth, which can vary species-to-species, and the tooth selected, with a single tooth from a high-crowned herbivore taking over a year to fully form and that of a small animal taking just weeks (Kohn and Cerling, 2002). For caribou and reindeer, estimations of enamel formation and mineralisation derive from studies of other deer species (e.g. Brown and Chapman, 1991a, 1991b). Based on these studies, and a recent case study using modern caribou (Britton et al., 2009), it is estimated that a full unworn *Rangifer* molar will represent between 6 and 9 months of growth, although consideration must be given to the influence that dental attrition and wear will have on this (Britton et al., 2009; Britton, 2010: 91). It must also be noted that sequential-sampling in this way does serve to homogenise enamel of each section into one datum point and, given the rapid mineralization time of these low-crowned teeth and the geometry of enamel growth, this may limit the resolution achievable.

The application of isotope analysis to mobility studies in archaeology and palaeoecology is made possible by the presumed relationship between the local environment (geology, climate) and the isotopic signatures of foods consumed by an individual. The use of strontium ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) and oxygen ( $\delta^{18}\text{O}$ ) isotope compositions from dental enamel (or ‘bioapatite’) are perhaps now the most established approaches in isotope palaeomobility studies, and have been utilised to investigate movements and migrations of both humans and animals in the past in a variety of geographical and temporal contexts (e.g. Britton et al., 2009; Eckardt et al., 2009; Evans et al., 2006; Henton et al., 2017; Hoppe, 2004; Valenzuela-Lamas et al., 2016). Strontium isotope approaches are based on the relationship between the characteristic strontium isotopic signatures of underlying lithologies and, by extension, location during tissue formation (see review in Bentley, 2006). Since strontium can substitute for calcium in hydroxyapatite, the mineral component of bone and tooth (McConnell, 1973) the  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios of faunal bioapatite can therefore be related to the region in which an organism sourced its food. This is particularly effective for wild herbivores, and the movements of archaeological and modern herbivores have been reconstructed using their strontium isotope composition (e.g. Britton

et al., 2009, 2011; Feranec et al., 2007; Hoppe et al., 1999).

The  $^{87}\text{Sr}/^{86}\text{Sr}$  of underlying bedrock varies primarily according to the age of the rock, as in geologically older rocks more of the radioactive mother isotope  $^{87}\text{Rb}$  has, over time, decayed to the radiogenic daughter isotope  $^{87}\text{Sr}$ . Therefore, older rocks have higher concentrations of  $^{87}\text{Sr}$ , and thus higher  $^{87}\text{Sr}/^{86}\text{Sr}$  (Capo et al., 1998). Although the geological age of rock is the primary influence on the  $^{87}\text{Sr}/^{86}\text{Sr}$ , other factors such as rock composition (i.e.  $^{87}\text{Rb}$  content) and mineralogy can affect this. Furthermore, while the isotopic ‘signature’ of soils or plants at any location (‘bioavailable’ strontium) is primarily influenced by the parent material/underlying lithology, other factors influence local bioavailable  $^{87}\text{Sr}/^{86}\text{Sr}$ , such as differential weathering processes, groundwater and precipitation, and atmospheric deposition such as sea spray (Bentley, 2006; Green et al., 2004; Montgomery et al., 2007; Vitousek et al., 1999). There are a number of approaches that can be used to estimate bioavailable  $^{87}\text{Sr}/^{86}\text{Sr}$  at a particular location, including the sampling of modern vegetation, soils, or near-ranging (micro-)fauna, or even the sampling of archaeological bone or dentine which (due to their susceptibility to diagenetic alteration) can provide an indication of the  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios of the surrounding burial environment at a site (Bentley and Knipper, 2005; Budd et al., 2000; Maurer et al., 2012; Price et al., 2002).

The  $\delta^{18}\text{O}$  values of precipitation are determined by climatic and environmental factors, primarily local temperature, but also altitude, humidity, and proximity to the coast (e.g. Dansgaard, 1964; Gat, 1980, 1996). In light of this, and the relationship between the  $\delta^{18}\text{O}$  values of mineralised tissues and that of ingested water (Bryant et al., 1994; Bryant and Froelich, 1995; Longinelli, 1984; Luz et al., 1984), oxygen isotopes can be utilised as a secondary indicator of geographical (climatic) origin in archaeological case studies. Furthermore, given that  $\delta^{18}\text{O}$  values of precipitation at mid- to high- latitudes vary seasonally,  $\delta^{18}\text{O}$  values can be used to ‘anchor’ strontium data within a seasonal context to further inform on seasonal movements and migrations (e.g. Britton et al., 2009; Julien et al., 2012; Pellegrini et al., 2008; Widga et al., 2010), while intra-tooth  $\delta^{18}\text{O}$  values (both the phosphate and carbonate components) can also be used to track local climatic conditions that occurred during tissue formation. The attenuation (‘dampening’) of the intra-tooth isotope signature, relative to the isotopic signatures of inputs (i.e. drinking water or food), occurs as a result of physiological factors, sampling strategy, and the nature, process and three-dimensional geometry of tooth mineralisation itself (Balasse, 2002; Kohn, 1996; Kohn, 2004; Kohn et al., 1996; Kohn and Cerling, 2002; Passey and Cerling, 2002). Additionally, it has been demonstrated that animal ranging behaviours, such as rapid, long-distance ‘thermo-stress’ migrations and altitudinal ranging, can further influence the nature of these attenuations (Britton et al., 2009; Tornero et al., 2016, 2017). It is on this basis that it has been suggested that attenuated oxygen isotope intra-tooth profiles in animals from areas of pronounced seasonal temperature variations (i.e. at mid- to high-latitudes) could indicate the scale and mode of seasonal ranging (near versus far ranging; rapid or gradual movements) undertaken by an animal (Britton et al., 2009).

### 3. The Nunalleq site

#### 3.1. Background

The Nunalleq site (59° 44' N, 161° 54' W) is a pre-contact Yup'ik village located in the Yukon-Kuskokwim Delta on the coast of the Bering Sea close to the village of Quinhagak, Western Alaska (Fig. 1). Investigations at the site followed a series of recent climatic changes that have resulted in, amongst things, an increase in unpredictable weather conditions in the region resulting in increased storms, a reduction in permafrost and, through this, coastal erosion which exposed the site in profile along the beach close to Quinhagak. Research and recovery efforts were subsequently undertaken at the request of the




Fig. 1. Map of Western Alaska showing the location of the Nunalleq site. (Source: adapted from Britton et al., 2013).

local Yup'ik descendant community. Between 2009 and 2015, excavations carried out by archaeologists from the University of Aberdeen uncovered a semi-subterranean multi-room sod house complex, which spans at least three different occupation phases between the 15th to 17th century AD (Britton et al., 2016; Ledger et al., 2016). The presence of discontinuous permafrost at the site has resulted in extremely-well preserved artefacts and ecofacts (including wooden tools, lithics, pottery, grass work, human hair, fur, animal bones and insects), a number of which have been the focus of recent studies (Britton et al., 2013, 2016; Farrell et al., 2014; Forbes et al., 2015; Ledger et al., 2016; McManus-Fry et al., 2016), contributing a wealth of data and building a wider picture of pre-contact life at Nunalleq. In a region that is sensitive to modern day climate change, and amongst communities that are suffering from its effects (Berkes and Jolly, 2001; Ford et al., 2006; Hansman, 2016; Pilkington, 2008). Nunalleq offers the chance to reconstruct and assess the impact that the most recent hemisphere-wide large scale climatic event (The Little Ice Age) had on the lives of the people of Nunalleq and the environment and animals around them, an especially important consideration given the modern-day context of similar environmental fragility.

#### 3.2. Regional climate, topography, and lithology

Nunalleq is located in the Yukon-Kuskokwim Delta region, an environment typified by a flat, waterlogged, tundra landscape. This area of Alaska is classified as subarctic Dfc in the Köppen climate classification, with a maritime influence from the Bering Sea. This climate type is defined as being a snow climate, fully humid with cool summers and cold winters (Kottek et al., 2006, after Köppen, 1900). Mean winter (January) temperatures in this area are between  $-13$  and  $-10$  °C (8.6 and 14°F), and between 11 and 18 °C (51.8 and 64.4°F) (mean) in summer (July) (Fleming et al., 2000). While only limited strontium isotope bioavailability data is available for Alaska, a recent geostatistical framework has been published for the state (based on measured lithological samples), creating a predictive model for both bedrock and




Fig. 2. Predicted  $^{87}\text{Sr}/^{86}\text{Sr}$  (median values) in bedrock in area surrounding the Nunalleq site and the modern Mulchatna herd range area (Source: adapted from Bataille et al., 2014: 6, Fig. 3B). Location of the Nunalleq site is shown.

river  $^{87}\text{Sr}/^{86}\text{Sr}$  (Bataille et al., 2014). Median values from the Bataille et al. (2014) study suggest  $^{87}\text{Sr}/^{86}\text{Sr}$  of local bedrock for the area around the site vary between 0.705 and 0.707 (see Fig. 2, adapted from Bataille et al., 2014: 6, Fig. 3B).

The site's main occupation period during the 15th to 17th century AD coincided with the Little Ice Age, a large scale climatic event that saw increased glaciation and generally cooler conditions. In the Northern hemisphere the period AD 1400 to 1900 is thought to have been around  $0.8\text{ }^{\circ}\text{C}$  ( $33.4^{\circ}\text{F}$ ) colder than the late 20th century, with the 17th and 19th century in particular being the coldest within this entire period (Mann et al., 1999; Mann, 2002). In most cases there appears to have been increased climatic variability, with intermittent periods of warmth punctuating the generally colder climate (Bradley and Jones, 1993; Pfister, 1995). Tree-ring palaeoclimate data from the Seward Peninsula (N.W. Alaska) confirms this, demonstrating cold conditions from the early to middle 1600s and again in the late 1700s to mid-1800s, with the most severe cold episode occurring around AD 1780 (D'Arrigo et al., 2005).


### 3.3. Caribou: past and present

The archaeological record at Nunalleq highlights the important role of caribou in the pre-contact Yup'ik world. Zooarchaeological evidence demonstrates that caribou carcasses were being brought to the site for processing (McManus-Fry, 2015: 160), skinning, consumption (meat and marrow) and for use as raw material in the osseous industry (antler and bone), with caribou remains making up approximately 22% of the faunal assemblage (total faunal NISP = 9852; Edouard Masson-

MacLean, pers. comm. 2017). While stable isotope evidence of human hair from Nunalleq suggests that salmon were likely the dominant protein source, diet at the site was mixed, incorporating terrestrial protein (likely caribou), fish (particularly anadromous species) and marine mammals (Britton et al., 2013, 2016). On the basis of isotopic mixing models, it has been estimated that caribou meat contributed approximately 9% of total dietary protein at Nunalleq (Britton et al., 2016), with domestic dogs at the site consuming proportionally a little more caribou meat than the human inhabitants of the site (McManus-Fry et al., 2016). Caribou are not just represented in the bioarchaeological record of Nunalleq, but are common in manufacturing-waste, artwork, and other finished pieces. Antler (debitage and artefacts) is a common find at the site, and examples of finished pieces featuring representations of caribou at the site include an antler engraved with a running caribou and hunting pendants in the shape of caribou hooves, attesting to the important relationship between this species and the inhabitants of Nunalleq.

Caribou remain an important subsistence species for the Yup'ik today, not only economically but culturally too, and like many animals they are traditionally considered sentient beings or 'non-human persons' (Fienup-Riordan, 2000; Simon-Sakurai, 2013). Contemporary changes in environmental conditions, however, appear to be affecting the availability of caribou, for example in Quinhagak where a lack of snow has resulted in fewer caribou in the area in recent years (Warren Jones, pers. comm. 2016). The modern caribou herd in this area are the Mulchatna, whose range includes the community of Quinhagak and the area immediately surrounding the Nunalleq archaeological site. The Mulchatna are a migratory herd of mountain caribou (*Rangifer tarandus*





**Fig. 3.** Map of modern day Mulchatna caribou herd seasonal ranges and movements, with location of the Nunalleq site shown.

(Source: adapted from image provided by Alaska Department of Fish and Game c/o Dominic Demma).

caribou) that undertake seasonal movements of up to 240 km (150 miles) over a range encompassing approximately 103,000 km<sup>2</sup> (approximately 40,000 m<sup>2</sup>) (Division of Wildlife Conservation, Alaska Department of Fish, and Game, 2016; and see Fig. 3). The area in which the herd currently calve is within their broader summer range, although split into two main areas. Both the summer range and the calving grounds incorporate areas of older lithology (including Mesozoic sedimentary and volcanic rock, and Paleozoic sedimentary, metamorphic, and igneous rock; see Beikman, 1980), with median bedrock values exceeding 0.709 and > 0.711 in the higher ground often favoured by mountain caribou during calving (Bataille et al., 2014: 6, Fig. 3B; and see Figs. 2 and 3 (this paper)).

#### 4. Materials and methods

Second (M2) and third (M3) molars were extracted from four archaeological caribou hemi-mandibles (individuals NUN- 1, -2, -3, -4, which correspond to bioarchaeological site sample identifiers B-248-534, -539, -541 and -540, respectively) and selected for strontium and oxygen (carbonate) isotope analysis. The material presented here come from two different phases of occupation at the site; the earlier phase III (NUN-1, 2, 3) and the later phase II (NUN-4). It should be noted that although material from the site has been radiocarbon dated, and stratigraphic relationships between these two phases are established, a more precise chronological model for the site has yet to be determined which currently limits diachronic investigations. Based on dental eruption and wear patterns it is estimated that these four individuals were aged between 2 and 3 years at death, with crown heights averaging between

10.3–15.5 mm (M2) and 10.6–15.7 mm (M3). No studies relating *Rangifer* tooth growth and development exist at present, however studies of other deer species can be used to gauge an estimation of the time period represented by a complete and unworn molar tooth (e.g. Brown and Chapman, 1991a, 1991b). It is estimated that *Rangifer* M2 and M3 teeth form sequentially with a M2 forming between < 3.5 and 9 months and a M3 between 9 and < 18 months (Brown and Chapman, 1991a). This estimation has been confirmed by previous sequential oxygen and strontium isotope studies of M2 and M3 enamel in modern caribou, although, in practice, periods represented in caribou/reindeer molars are slightly reduced due to wear that removes some enamel containing a few months of isotope data at the crown, so that the time period represented is thought to be approximately 1 year (Britton et al., 2009; Britton, 2010). While these modern studies have highlighted correlations between intra-tooth strontium and oxygen isotope profiles in *Rangifer* molars and changes in these isotope systems and migratory behaviours, it should be noted that uncertainties do surround the phasing and periodicity of mineralisation in *Rangifer* enamel and thus their intra-tooth isotopic profiles. For example, it cannot be unequivocally assumed that the strontium and oxygen isotopic signals are incorporated into tooth enamel synchronously. While this does not limit current applications (given that most methods of intra-tooth sampling tend to target multiple growth layers, somewhat homogenising isotopic ‘signals’ from neighbouring portions of enamel), further experimental studies would be needed in order to investigate this, and other aspects of *Rangifer* tooth development and mineralisation, in order to facilitate the resolution achievable in incremental enamel isotope studies were more fine-grained sampling approaches employed (e.g. laser ablation).

#### 4.1. Sampling

Molars were extracted from the bone using a stainless-steel circular saw, brush-cleaned with deionised (DI) water, and then left to dry overnight. Whole teeth were then mechanically abraded using a tungsten carbide burr (NTI-Kahla, Germany) to remove surficial enamel and any adhering material, ultrasonicated in DI water, and dried. Intra-tooth sampling was performed using flexible, stainless steel diamond-coated superfine circular bits (NTI-Kahla, Germany), with drill bits cleaned between each sample using a weak nitric acid (< 0.3 M) to dissolve any adhering enamel, rinsed with DI water, ultrasonicated for 5 minutes in DI water and left to dry before re-use.

Enamel from the buccal face of the anterior loph of each tooth was selected for sampling, following the method detailed in Britton et al. (2009). After cutting and cleaning, enamel faces were measured, marked, and cut as horizontal sequential samples (perpendicular to the occlusal surface) at ~1.5 mm intervals. Sections were given sequential numerical sample assignments, starting from the enamel-root junction (ERJ) to the occlusal surface (M2-E1, M2-E2, M2-E3, etc.), and sections were split longitudinally, with ~5 mg of enamel reserved for each isotope analyses. A sample of (root) dentine was retained for bulk strontium isotope analysis.

#### 4.2. Strontium

Strontium was extracted using the method from Deniel and Pin (2001) with modifications outlined in Copeland et al. (2008), at the clean laboratory facility, Department of Archaeology, Memorial University, Newfoundland (Canada). Clean enamel samples (~5 mg) were rinsed once with ~200 µl of ultra-pure acetone and left to dry completely. Samples were then dissolved in closed 3 ml Savillex vials (Minnetonka, MN, USA) with 1.5 ml 8 M HNO<sub>3</sub> (single distilled Reagent grade, Sigma Aldrich) at 70 °C. After samples were fully digested 0.5 ml HNO<sub>3</sub> was removed to determine Sr concentration. The remaining solution (1 ml) was loaded into pre-conditioned micro columns (after Charlier et al., 2006) containing Sr-spec resin (EiChrom, Darian, IL, USA). After three successive washes using 1 ml of 8 M HNO<sub>3</sub>, strontium was eluted from the resin in 2 ml deionized water and then acidified with the addition of 75 µl of 8 M HNO<sub>3</sub>, ready for analysis of <sup>87</sup>Sr/<sup>86</sup>Sr ratios using a Neptune Multi-Collector Inductively Coupled Plasma Mass Spectrometer (MC-ICP-MS, Thermo Fisher Scientific, Bremen, Germany). A 200 ppb solution of Sr isotope standard SRM 987 (strontium carbonate dissolved in 0.3 N HNO<sub>3</sub>, National Institute of Standards and Technology, Gaithersburg, MD, USA) was analysed between every five samples for quality control purposes. Using the average <sup>87</sup>Sr/<sup>86</sup>Sr value for all the SRM 987 collected in the analytical session, all data was externally corrected to the SRM 987 value of <sup>87</sup>Sr/<sup>86</sup>Sr = 0.710240 (Johnson and Fridrich, 1990). The facility's long-term average for SRM 987 from 2014–2016 was 0.710268 ± 0.000028 (2SD, n = 338).

#### 4.3. Oxygen

Enamel was prepared for carbonate oxygen isotope analysis following the methods outlined in Koch et al. (1997), with modifications detailed in Britton et al. (2009: 1166). Samples were powdered using an agate pestle and agitated in 30% H<sub>2</sub>O<sub>2</sub> (40 µl/1 mg) at room temperature for 24 hours. Samples were then rinsed five times with DI water and dried for 48 hours at 40 °C. Samples were measured for carbonate oxygen isotope ratios at Johannes Gutenberg Universität Mainz, Germany and Iso-Analytical Ltd., Sandbach, Cheshire, UK. In Mainz, ~1 mg aliquots of the dried pre-treated powders were analysed using a Thermo Gasbench II coupled via a ConFlo II to a ThermoFinnigan MAT 253 gas mass spectrometer. At Iso-Analytical Ltd. sample sizes were larger (3.8 mg to 8.4 mg), and weighed into vials that were then flushed with 99.995% helium. Phosphoric acid was then added to vials and left overnight to react to allow for complete conversion of carbonate to

CO<sub>2</sub>. The CO<sub>2</sub> gas was analysed using CF-IRMS (Europa Scientific 20-20 IRMS). In both laboratories, values were normalised to in-house and international standards (NBS-18, NBS-19, Carrara Marble), and analytical error for δ<sup>18</sup>O<sub>CO<sub>2</sub></sub> measurements was ± 0.2 ‰ (1 SD) and ± 0.1 ‰ (1 SD) (or better) at Iso-Analytical Ltd. and Mainz respectively, based on repeat analyses of those standards. This analytical error was also confirmed by the repeated analysis of an internal enamel reference material (horse tooth enamel) in analytical runs. All δ<sup>18</sup>O<sub>CO<sub>2</sub></sub> values are reported here relative to the international reference material VSMOW.

### 5. Results and discussion

The results of the isotopic measurements are summarised in Table 1. Sequential strontium and oxygen isotope data from the four individuals are presented in Figs. 4 and 5, respectively. The values have been plotted aligning the most recently formed parts of each tooth (i.e. M2-E1, M3-E1) to allow inter-individual comparison. ‘Anchoring’ and plotting the data from the ERJ in this way ensures cross comparability and is a common approach in incremental studies when using variable sampling increments (e.g. Balasse et al., 2012; Buchan et al., 2016) or fixed ones (e.g. Britton et al., 2009, 2011; this study). Fig. 6a–d show the individual strontium and oxygen isotope profiles from each of the individuals, allowing a direct comparison of the intra- and inter-tooth <sup>87</sup>Sr/<sup>86</sup>Sr and δ<sup>18</sup>O<sub>CO<sub>2</sub></sub> trends.

The total isotopic range for all intra-tooth data from all individuals is 0.7057 ± 0.0003 (<sup>87</sup>Sr/<sup>86</sup>Sr), with the lowest individual mean of 0.7062 ± 0.0003 (1SD, n = 14) and highest individual mean of 0.7067 ± 0.0001 (1SD, n = 12), and from 17.4 ‰ to 21.6 ‰, with a lowest individual mean of 18.2 ‰ ± 0.7(1SD, n = 11) and highest individual mean of 20.4 ± 0.6 ‰ (1SD, n = 10) (δ<sup>18</sup>O<sub>CO<sub>2</sub></sub> VSMOW). The strontium isotope values determined from the dentine (n = 8) are higher than the values measured in the enamel. These values ranged from 0.7064 to 0.7077, with a mean of 0.7071 ± 0.0004 (1SD). Given the coastal location of the site, it is possible that seawater strontium (as sea spray) has influenced the depositional environment and thus the dentine values (Whipkey et al., 2000).

#### 5.1. Sequential strontium isotope data

The strontium isotope profiles for all four individuals display the same overall, cyclical trend from lower values to higher to lower again, suggesting some seasonal movement (Fig. 4). All individuals demonstrate intra-tooth <sup>87</sup>Sr/<sup>86</sup>Sr variation in the third or fourth decimal place and display a similar trend in their <sup>87</sup>Sr/<sup>86</sup>Sr, from lower values in the earliest forming parts (0.7063 ± 0.0002, 1SD, at point M2-E6) to more elevated values (0.7067 ± 0.0002, 1SD, at point M2-E1) and a return to lower values (0.7062 ± 0.0004, 1SD, at point M3-E1) over a time period thought to represent approximately one year of life (Britton et al., 2009).

The relatively low intra-tooth variation/range of values for individuals NUN-1 and NUN-2 (0.7061 to 0.7065 and 0.7066 to 0.7068, respectively) suggests minimal movement over young, relatively isotopically homogenous lithologies. The intra-tooth strontium isotope profiles of NUN-3 and NUN-4 demonstrate slightly more variation, with values ranging from 0.7063 to 0.7071 in NUN-3 and 0.7057 to 0.7065 in NUN-4 across all sections taken in both teeth, perhaps suggesting movement over slightly more isotopically- (i.e. geologically) varied terrain. However, inter-individual variation is like that determined in modern herds (Britton et al., 2009), suggesting similar migration-histories. Indeed, the inter-individual similarities in all four caribou in the earliest forming parts of the M2 (e.g. ± 0.0002 at point M2-6) indicate spatial proximity in the first few months of life. This would likely correspond with the latter phases of infancy on and around the calving grounds and summer range of these individuals, as has been observed in experimental studies on caribou from modern caribou herds of known movement (Britton et al., 2009; Britton, 2010). Further similarities in


**Table 1**

Summary of isotopic results ( $\delta^{18}\text{O}_{\text{CO}_3 \text{ VSMOW}}$  and  $^{87}\text{Sr}/^{86}\text{Sr}$ ) from sequentially-sampled enamel (E) of second and third molars of four caribou from Nunalleq. Sections were given numerical assignments commencing from the enamel-root junction (ERJ) to the occlusal surface. Also presented are the bulk dental (D)  $^{87}\text{Sr}/^{86}\text{Sr}$  data for each individual.

Individual	Sample	$\delta^{18}\text{O}_{\text{CO}_3 \text{ VSMOW}}$	$^{87}\text{Sr}/^{86}\text{Sr}$	Individual	Sample	$\delta^{18}\text{O}_{\text{CO}_3 \text{ VSMOW}}$	$^{87}\text{Sr}/^{86}\text{Sr}$
NUN-1 (B-248-534)	M2-E9	No data	0.7061	NUN-2 (B-248-539)	M2-E6	No data	0.7066
	M2-E8	No data	0.7061		M2-E5	19.6	0.7066
	M2-E7	19.8	0.7061		M2-E4	18.9	0.7066
	M2-E6	19.6	0.7061		M2-E3	19.3	0.7067
	M2-E5	18.6	0.7061		M2-E2	18.4	0.7067
	M2-E4	18.7	0.7063		M2-E1	18.5	0.7068
	M2-E3	18.5	0.7064		M3-E6	18.5	0.7069
	M2-E2	17.6	0.7063		M3-E5	18.7	0.7069
	M2-E1	No data	0.7064		M3-E4	19.0	0.7068
	M3-E10	17.9	0.7065		M3-E3	19.2	0.7068
	M3-E9	17.8	0.7065		M3-E2	19.4	0.7067
	M3-E8	17.6	0.7065		M3-E1	No data	0.7066
	M3-E7	17.7	0.7065		M2-D	No data	0.7073
	M3-E6	18.1	0.7065		M3-D	No data	0.7069
	M3-E5	18.8	0.7064				
	M3-E4	19.6	0.7064				
	M3-E3	20.0	0.7064				
	M3-E2	20.6	0.7063				
	M3-E1	20.5	0.7062				
	M2-D	No data	0.7077				
M3-D	No data	0.7076					
NUN-3 (B-248-541)	M2-E7	21.63	0.7064	NUN-4 (B-248-540)	M2-E7	17.8	0.7062
	M2-E6	20.78	0.7064		M2-E6	17.7	0.7062
	M2-E5	20.44	0.7064		M2-E5	17.6	0.7063
	M2-E4	20.09	0.7066		M2-E4	17.4	0.7064
	M2-E3	19.97	0.7067		M2-E3	17.6	0.7066
	M2-E2	19.58	0.7068		M2-E2	17.8	0.7066
	M2-E1	No data	0.7069		M2-E1	No data	0.7065
	M3-E6	19.81	0.7071		M3-E7	18.4	0.7064
	M3-E5	19.91	0.7070		M3-E6	19.0	0.7062
	M3-E4	20.17	0.7068		M3-E5	19.1	0.7061
	M3-E3	21.02	0.7066		M3-E4	19.0	0.7060
	M3-E2	20.94	0.7065		M3-E3	18.7	0.7059
	M3-E1	No data	0.7063		M3-E2	18.9	0.7058
M2-D	No data	0.7064	M3-E1	No data	0.7057		
M3-D	No data	0.7068	M2-D	No data	0.7073		
			M3-D	No data	0.7069		

later-forming portions of both teeth further demonstrate this likely spatial proximity throughout much of the first year of life and indicate that these four animals were moving over similar lithologies, likely within the same broad geographical area. In all individuals, the  $^{87}\text{Sr}/^{86}\text{Sr}$  closest to that of the mean 'local' value, as indicated by dentine values ( $0.7071 \pm 0.0004$ , 1SD,  $n = 8$ ), occur in the latest-forming parts of the M2 and earliest forming parts of the M3. This may indicate these caribou were near Nunalleq during this period, perhaps

giving the inhabitants of Nunalleq their hunting opportunity. This hypothesis is not without merit; zooarchaeological analysis of the age at death of caribou from Nunalleq point to the likelihood of a winter hunt, although hunts may have occurred at other times of the year too (McManus-Fry, 2015). The caribou 'search and harvest areas' (hunting areas) used by the inhabitants of Quinhagak today are approximately between 60 (north and southwards) and 120 (eastwards) km (37 and 75 miles) away, so it could be likely that the traditional hunting



**Fig. 4.** Plot of all strontium isotope data ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) from intra-tooth sections from M2 and M3 enamel of each of the four Nunalleq caribou. Closed symbols are Phase III individuals (triangle = NUN-1, diamond = NUN-2, circle = NUN-3) and open symbols are the Phase II individual (square = NUN-4). The mean dentinal value ( $0.7071 \pm 0.0004$  1SD;  $n = 8$ ) is represented by the dashed line. Analytical error is within the symbol markers.




Fig. 5. Plot of all  $\delta^{18}\text{O}_{\text{Co}3}$  data from intra-tooth sections from enamel from second and third molars of four caribou from Nunalleq. Closed symbols are Phase III individuals (triangle = NUN-1, diamond = NUN-2, circle = NUN-3) and open symbols are the Phase II individual (square = NUN-4).

grounds were of a similar distance away from the site of Nunalleq (Simon, 2016: 57).

The  $^{87}\text{Sr}/^{86}\text{Sr}$  from all individuals are within the range of expected values for the area surrounding the site, which is predicted to be between 0.705 and 0.707 based on a predicted geospatial model for  $^{87}\text{Sr}/^{86}\text{Sr}$  in Alaskan rocks (Bataille et al., 2014: 6, Fig. 3B), and are also similar to the  $^{87}\text{Sr}/^{86}\text{Sr}$  of dentine measured from the site (presumed to reflect locally bioavailable Sr), which range from 0.7064 to 0.7077, with a mean of  $0.7071 \pm 0.0004$  (1SD,  $n = 8$ ; see Table 1). Given that the present-day Mulchatna Herd in this region calve and spend the summer months in areas which incorporate older lithologies (with expected higher  $^{87}\text{Sr}/^{86}\text{Sr}$ ; see Figs. 2 and 3, and Bataille et al., 2014: 6:

Fig. 3B), it seems unlikely that the Nunalleq caribou utilised the same calving grounds and broader summer range as the Mulchatna caribou use today. This is especially interesting given that calving ground location in particular is thought to be quite stable (Ferguson and Elkie, 2004; Russell et al., 1993). Given the more lithologically varied terrain encompassed in the total range of the Mulchatna herd today, it is unlikely that the geographical range used by the Nunalleq caribou was the same as modern populations in the region or perhaps only incorporated part of what is used today. Given the lack of bioavailability data for the broader region, and the relative homogeneity of lithology to the northeast of the site, the precise location of the calving and summer range areas used by these archaeological caribou cannot be determined. It can




Fig. 6. Plots a–d demonstrate intra-tooth isotope values of  $^{87}\text{Sr}/^{86}\text{Sr}$  (solid circles) and  $\delta^{18}\text{O}_{\text{Co}3}$  (open triangles) for M2 and M3s of each of the caribou analysed. Plot a, b and c represent the phase III individuals NUN-1, -2 and -3, respectively. Plot d represents the phase II individual NUN-4. Analytical error for strontium measurements is within the symbol markers.



be noted that there is a large swathe of Quaternary lithology (sedimentary) extending northwards from Nunalleq to the Norton Sound area. Beyond this are areas of older rock (e.g. Mesozoic sedimentary rock, Upper Paleozoic metamorphic and igneous rock; see Fig. 2). It is perhaps likely then that these caribou were born in, and ranged throughout, this relatively ‘penned-in’ area of Quaternary lithology.

Recent large shifts in the calving areas used by the Mulchatna herd have been observed during annual survey work and are thought to have resulted from increased predation, short-term weather affects as well as longer-term climate change (Dominic Demma, pers. comm. 2016). Similar abandonments and shifts in calving areas have been observed for other caribou herds in Canada and Alaska (e.g. Abraham et al., 2012; Cameron et al., 1992). In Greenland, an association has also been demonstrated between caribou population size and climate change, and it has been shown that caribou herds undergo ‘boom-bust’ cycles, with herd sizes expanding and contracting enormously in a relatively periodic fashion and in relation to climatic shifts in relatively recent history (Meldgaard, 1986: 59–64). Similar population fluctuations, albeit not regular cycles, in caribou herds have been observed in Alaska (e.g. Bergerud et al., 1984; Skoog, 1968) and Canada (e.g. Couturier et al., 1990) and more recent studies, using long-term satellite and radiocollar observation, have identified similar trends in caribou herd population-size peaks and troughs (e.g. Abraham et al., 2012; Hegel and Russell, 2013; Ruttan, 2012). The reasons for these, apparently cyclical, fluctuations in herd numbers, and the trigger behind such extreme ‘boom-bust’ cycles, may be (local and global) climatic variations, forage availability and changes in predation risks (Gunn, 2003; Meldgaard, 1986; McNeil et al., 2005; Weladji et al., 2002). The relationship between fluctuations in overall herd size and range use in past caribou populations requires further research, although how past herd sizes could be determined remains unclear. Additionally, further investigation into the implications of such changes in caribou behaviours for contemporary human groups would be of great interest both to archaeologists and present-day Indigenous communities. Future work, including the integration of these isotopic data with site-specific diachronic human stable isotope palaeodietary data, more detailed zooarchaeological analyses and material culture changes (particularly of hunting implements), will undoubtedly prove illuminating. The integration of ethno-historical accounts and oral histories into such studies would also prove valuable.

It is also possible that, rather than documenting the *changing* behaviour of the Mulchatna, the strontium data from Nunalleq instead evidence the behaviour of another (now extinct) population of caribou. Indeed, there was a historically-documented population of caribou (Bering Sea Coast Herds) that were abundant along the lowland areas of the Bering Sea coast until the early 1900s when they disappeared entirely (Murie, 1935, after Nelson, 1888). Although not a distinct herd, these caribou appear to have undertaken a general north-south migration from the Norton Sound area to Kuskokwim Bay (Murie, 1935: 61). Such a migratory route would have incorporated the vicinity of the site.

### 5.2. Sequential oxygen isotope data

Oxygen isotope compositions for the four individuals also display a similar trend through time, from higher  $\delta^{18}\text{O}_{\text{CO}_3}$  values, to lower values, before a return to higher  $\delta^{18}\text{O}_{\text{CO}_3}$  values again (Fig. 5). These fluctuations in  $\delta^{18}\text{O}_{\text{CO}_3}$  values are likely a result of seasonal variations in temperature during the period of tooth growth. The higher  $\delta^{18}\text{O}_{\text{CO}_3}$  isotope values correspond to summer, the lower values to winter.

Intra-tooth oxygen values in individuals NUN-1 and NUN-2 are similar in terms of overall trend and measured values. These values vary from  $19.1 \pm 0.7 \text{ ‰}$  (1SD) at point M2-5 to  $18.0 \pm 0.6 \text{ ‰}$  (1SD) at point M2-E2 to  $20.0 \pm 0.8 \text{ ‰}$  (1SD) at point M3-E2. This indicates that, after being born in the same season (albeit very likely across different years), the  $\delta^{18}\text{O}_{\text{CO}_3}$  values of water ingested during the first year of life (either directly, or via plants consumed) was similar for

these different animals. The same trend is also visible in individual NUN-3; however, the measured values are slightly higher. This could be due to intra or inter-annual climatic variations, or even due to differences in feeding behaviours and the water physiology of plants consumed.

These three animals, from phase III, demonstrate greater intra-tooth variation (0.4–1.0‰) than inter-tooth variation (i.e. means for M2 and M3 measurements are similar in the same animal). The intra-tooth values fluctuate in both teeth, with summer peaks and winter troughs, demonstrating the seasonality of birth and the period of tooth development. Measurements in the later-forming parts of the M2 and the earliest forming parts of the M3 are similar across all three individuals; NUN-1 (17.6‰ and 17.9‰ at points M2-E2 and M3-E10 respectively), NUN-2 (18.5‰ at points M2-E1 and M3-E6), NUN-3 (19.6‰ and 19.8‰ at points M2-E2 and M3-E6 respectively).

Individual NUN-4 displays a slightly different trend in  $\delta^{18}\text{O}_{\text{CO}_3}$  values. The values vary from 17.8‰ at M2-E7, to 17.4‰ at M2-E4, to 18.9‰ at M3-E2. These absolute values, especially within the M2, are lower ( $\sim 1.3$ – $2.8 \text{ ‰}$ ) compared to the other three individuals’ values at equivalent points in tooth development. Given that the  $^{87}\text{Sr}/^{86}\text{Sr}$  in the M2 from this same individual indicates spatial proximity with the other three individuals at/around the time of birth, it is perhaps likely that the difference in absolute values represent a difference in seasonal (spring/summer) temperatures between these phases/time periods. However, NUN-4 also shows little intra-tooth variation (0.6‰) but more inter-tooth variation ( $\sim 1.2 \text{ ‰}$ ), which may reflect a substantial shift in location or environment between the period of M2 and M3 formation, and perhaps suggests this individual – while traversing similar lithological units to the other individuals studied – may have undertaken a seasonal migratory behaviour of a different nature from the other individuals (e.g. varying in distance, north-south directionality, and/or altitude).

### 5.3. Inter-individual variability in $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{18}\text{O}_{\text{CO}_3}$

Particularly in  $\delta^{18}\text{O}_{\text{CO}_3}$ , individual NUN-4 demonstrates a different isotopic profile compared to the other three caribou (see Fig. 6a–d). The intra-tooth variation in  $\delta^{18}\text{O}_{\text{CO}_3}$  for NUN-1, 2 and 3 – in combination with the  $^{87}\text{Sr}/^{86}\text{Sr}$  variation – suggests seasonal movement but over a restricted range. Conversely, the lower  $\delta^{18}\text{O}_{\text{CO}_3}$  intra-tooth variation observed in NUN-4 is more comparable to modern migratory caribou undertaking north-south trajectory migrations (Britton et al., 2009). It is possible therefore that this animal was undertaking more significant seasonal movements compared to the three caribou from Phase III. The lack of intra-tooth  $\delta^{18}\text{O}_{\text{CO}_3}$  variation is also perhaps representative of this caribou being a long-distance thermo-stress migrant since the observed ‘dampening’ in the intra-tooth oxygen isotope values is likely further influenced by rapid, seasonal migrations (Britton et al., 2009). An alternative hypothesis which would similarly explain the greater attenuation in  $\delta^{18}\text{O}_{\text{CO}_3}$  intra-tooth values is that this caribou was born at a higher elevation and ranged altitudinally, since this type of mobility can also serve to reduce the annual amplitude of isotopic variation resulting in ‘dampened’ values (Tornero et al., 2016, 2017). Indeed, the nearby Kuskokwim and Kilbuck mountain ranges are known to have been used by caribou herds both in the historical past and in more recent times (Simon, 2016). Additionally, it is possible that this individual was of a different sex compared to the other three. Sexual segregation and differences in habitat use have been observed in other modern caribou herds (e.g. Jakimchuk et al., 1987), which could potentially result in possible differences in the timing, scope, or distance travelled during migrations.

This observed inter-individual variability, particularly in  $\delta^{18}\text{O}_{\text{CO}_3}$ , may point to between-phase, and thus between time period, differences in the isotopic profiles (and perhaps the climate and/or ranging behaviours) of these caribou although, with only a single individual from phase II, estimation of behavioural differences of caribou between the

two occupation phases cannot be further elaborated upon. Strontium and oxygen isotope measurements from individuals from the modern herd in the region today would aid in better understanding intra-group isotopic variability across this broader area, although unfortunately none were available for this study.

## 6. Conclusions

This study has provided the first isotopic evidence for caribou seasonal movements in pre-contact Alaska, and in this region during the Little Ice Age. Here intra-tooth  $^{87}\text{Sr}/^{86}\text{Sr}$  and  $\delta^{18}\text{O}_{\text{CO}_3}$  data from caribou bioapatite from the site of Nunalleq indicate that these four animals were undertaking seasonal migrations, albeit perhaps limited in nature. In all individuals, the  $^{87}\text{Sr}/^{86}\text{Sr}$  closest to the locally bioavailable  $^{87}\text{Sr}/^{86}\text{Sr}$  occur in winter and may indicate these caribou were near Nunalleq during this period, perhaps giving the inhabitants of Nunalleq their hunting opportunity. Possible inter-individual differences in  $\delta^{18}\text{O}_{\text{CO}_3}$  have been discussed, which may be related to differences in climate and/or ranging behaviours between phases of site occupation. Further work to establish the nature of the change in climate during the site's occupation is ongoing and may provide the opportunity to correlate any climatic or environmental shifts with the possible changes in behaviour observed here. The comparison of the  $^{87}\text{Sr}/^{86}\text{Sr}$  data from all archaeological individuals with values expected for modern caribou in the region (given the known locations of the local present-day herd's range) may indicate a shift in the location of calving grounds/summer range, and possibly the size of overall range, between the 15th to 17th century AD and the present day. The reasons for this apparent shift can only be speculated upon, however, it may be that the period of relatively rapid climatic warming which has occurred in this region since the end of the Little Ice Age may have played a part. Additionally, it is possible that these Nunalleq caribou represent members of an extinct population that historically roamed along the Bering Sea coast. The analysis of additional *Rangifer* remains from historical, pre-contact and prehistoric contexts in the region and more broadly across the circumpolar North is clearly needed to better investigate the impact of climate change on caribou/reindeer range use and ecological behaviours. Ideally, however, this should be coupled with the analysis of modern environmental samples as part of broader bioavailability studies and with the experimental analysis of modern animals, analysed alongside accompanying telemetry and satellite survey information.

## Acknowledgements

This work was funded by an Arts and Humanities Research Council (AH/K006029/1) grant awarded to Rick Knecht, Kate Britton and Charlotta Hillerdal (Aberdeen). The onsite collection of samples was carried out by staff and students from the University of Aberdeen, volunteer excavators and the residents of Quinhagak. We also thank the Qanirtuuq Incorporated, Quinhagak, Alaska, and the people of Quinhagak for logistical and planning support for fieldwork and sampling permissions. Special thanks for Warren Jones and Qanirtuuq Incorporated (especially Michael Smith and Lynn Church), and to all Nunalleq project team members, particularly Edouard Masson-MacLean and Paul Ledger (Aberdeen). Thanks also to Alison Harris (Stockholm/York), Rebecca Lam and Sherri Strong (Memorial University) and Michael Maus (Johannes-Gutenberg Universität, Mainz) for methodological and technical assistance. Thank you to Dominic Demma (Alaska Department of Fish and Game) for providing information on the Mulchatna herd, to Edouard Masson-MacLean for comments on an earlier draft of this manuscript, and to the guest editors for inviting us to contribute to this special issue. Finally, the authors would like to thank the reviewers for their valuable and constructive comments on this manuscript.

## References

- Abraham, K.F., Pond, B.A., Tully, S.M., Trim, V., Hedman, D., Chenier, C., Racey, G.D., 2012. Recent changes in summer distribution and numbers of migratory caribou on the southern Hudson Bay coast. *Rangifer* 32, 269–276. <http://dx.doi.org/10.7557/2.32.2.2275>.
- Balasse, M., 2002. Reconstructing dietary and environmental history from enamel isotopic analysis: time resolution of intra-tooth sequential sampling. *Int. J. Osteoarchaeol.* 12, 155–165. <http://dx.doi.org/10.1002/oa.601>.
- Balasse, M., 2003. Potential biases in sampling design and interpretation of intra-tooth isotope analysis. *Int. J. Osteoarchaeol.* 13, 3–10. <http://dx.doi.org/10.1002/oa.656>.
- Balasse, M., Tresset, A., 2002. Early weaning of Neolithic domestic cattle (Bercy, France) revealed by intra-tooth variation in nitrogen isotope ratios. *J. Archaeol. Sci.* 29, 853–859. <http://dx.doi.org/10.1006/jasc.2001.0725>.
- Balasse, M., Smith, A.B., Ambrose, S.H., Leigh, S.R., 2003. Determining sheep birth seasonality by analysis of tooth enamel oxygen isotope ratios: the late Stone Age site of Kasteelberg (South Africa). *J. Archaeol. Sci.* 30, 205–215 (doi: 10.1006/jasc.2002.0833).
- Balasse, M., Tresset, A., Ambrose, S.H., 2006. Stable isotope evidence ( $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$ ) for winter feeding on seaweed by Neolithic sheep of Scotland. *J. Zool.* 270, 170–176. <http://dx.doi.org/10.1111/j.1469-7998.2006.00104.x>.
- Balasse, M., Boury, L., Ughetto-Monfrin, J., Tresset, A., 2012. Stable isotope insights ( $\delta^{18}\text{O}$ ,  $\delta^{13}\text{C}$ ) into cattle and sheep husbandry at Bercy (Paris, France, 4th millennium BC): birth seasonality and winter leaf foddering. *Environ. Archaeol.* 17, 29–44. <http://dx.doi.org/10.1179/1461410312Z.0000000003>.
- Bataille, C.P., Brennan, S.R., Hartmann, J., Moosdorf, N., Wooller, M.J., Bowen, G.J., 2014. A geostatistical framework for predicting variations in strontium concentrations and isotope ratios in Alaskan rivers. *Chem. Geol.* 389, 1–15. <http://dx.doi.org/10.1016/j.chemgeo.2014.08.030>.
- Beikman, H.M., 1980. Geologic Map of Alaska. U.S. Geological Survey (Map SG0002-1T and 2T. 1: 2 500 000).
- Bendrey, R., Vella, D., Zazzo, A., Balasse, M., Lepetz, S., 2015. Exponentially decreasing tooth growth rate in horse teeth: implications for isotopic analyses. *Archaeometry* 57, 1104–1124. <http://dx.doi.org/10.1111/arc.12151>.
- Bentley, R.A., 2006. Strontium isotopes from the earth to the archaeological skeleton: a review. *J. Archaeol. Method. Th.* 13, 135–187. <http://dx.doi.org/10.1007/s10816-006-9009-x>.
- Bentley, R.A., Knipper, C., 2005. Geographical patterns in biologically available strontium, carbon, and oxygen isotope signatures in prehistoric SW Germany. *Archaeometry* 47, 629–644.
- Bergerud, A.T., 1988. Caribou, wolves, and man. *Trends Ecol. Evol.* 3, 68–72.
- Bergerud, A.T., Jakimchuk, R.D., Carruthers, D.R., 1984. The buffalo of the north: caribou (*Rangifer tarandus*) and human developments. *Arctic* 37, 7–22.
- Berkes, F., Jolly, D., 2001. Adapting to climate change: social-ecological resilience in a Canadian western arctic community. *Conserv. Ecol.* 5, 1–15.
- Blois, J.L., Hadly, E.A., 2009. Mammalian response to Cenozoic climatic change. *Annu. Rev. Earth Planet. Sci.* 37, 181–208. <http://dx.doi.org/10.1146/annurev.earth.031208.100055>.
- Bradley, R.S., Jones, P.D., 1993. “Little Ice Age” summer temperature variations: their nature and relevance to recent global warming trends. *The Holocene* 3, 367–376.
- Britton, K., 2010. Multi-isotope Analysis and the Reconstruction of Prey Species Palaeomigrations and Palaeoecology. PhD Thesis. Durham University.
- Britton, K. in press. Prey species movements and migrations in ecocultural landscapes: reconstructing late Pleistocene herbivore seasonal spatial behaviours, in: Pilaar-Birch, S. (Ed.), *Multi-Species Archaeology*. Routledge, London, pp. 1–24.
- Britton, K., Grimes, V., Dau, J., Richards, M.P., 2009. Reconstructing faunal migrations using intra-tooth sampling and strontium and oxygen isotope analyses: a case study of modern caribou (*Rangifer tarandus granti*). *J. Archaeol. Sci.* 36, 1163–1172. <http://dx.doi.org/10.1016/j.jas.2009.01.003>.
- Britton, K., Grimes, V., Niven, L., Steele, T.E., McPherron, S., Soressi, M., Kelly, T.E., Jaubert, J., Hublin, J.-J., Richards, M.P., 2011. Strontium isotope evidence for migration in late Pleistocene *Rangifer*: implications for Neanderthal hunting strategies at the Middle Palaeolithic site of Jonzac, France. *J. Hum. Evol.* 61, 176–185. <http://dx.doi.org/10.1016/j.jhevol.2011.03.004>.
- Britton, K., Knecht, R., Nehlich, O., Hillerdal, C., Davis, R.S., Richards, M.P., 2013. Maritime adaptations and dietary variation in prehistoric Western Alaska: stable isotope analysis of permafrost-preserved human hair: diet in prehistoric Western Alaska. *Am. J. Phys. Anthropol.* 151, 448–461. <http://dx.doi.org/10.1002/ajpa.22284>.
- Britton, K., Fuller, B.T., Tütken, T., Mays, S., Richards, M.P., 2015. Oxygen isotope analysis of human bone phosphate evidences weaning age in archaeological populations. *Am. J. Phys. Anthropol.* 157, 226–241. <http://dx.doi.org/10.1002/ajpa.22704>.
- Britton, K., McManus-Fry, E., Nehlich, O., Richards, M., Ledger, P.M., Knecht, R., 2016. Stable carbon, nitrogen and sulphur isotope analysis of permafrost preserved human hair from rescue excavations (2009, 2010) at the precontact site of Nunalleq, Alaska. *J. Archaeol. Sci. Rep.* <http://dx.doi.org/10.1016/j.jasrep.2016.04.015>.
- Brotton, J., Wall, G., 1997. Climate change and the Bathurst caribou herd in the north-west territories, Canada. *Clim. Chang.* 35, 35–52. <http://dx.doi.org/10.1023/A:1005313315265>.
- Bryant, J.D., Froelich, P.N., 1995. A model of oxygen isotope fractionation in body water of large mammals. *Geochim. Cosmochim. Acta* 59, 4523–4537. [http://dx.doi.org/10.1016/0016-7037\(95\)00250-4](http://dx.doi.org/10.1016/0016-7037(95)00250-4).
- Bryant, J.D., Luz, B., Froelich, P.N., 1994. Oxygen isotopic composition of fossil horse tooth phosphate as a record of continental paleoclimate. *Palaeogeogr.*

- Palaeoclimatol. Palaeoecol. 107, 303–316. [http://dx.doi.org/10.1016/0031-0182\(94\)90102-3](http://dx.doi.org/10.1016/0031-0182(94)90102-3).
- Brown, W.A.B., Chapman, N.G., 1991a. The dentition of red deer (*Cervus elaphus*): a scoring scheme to assess age from wear of the permanent molariform teeth. *J. Zool.* 224, 519–536.
- Brown, W.A.B., Chapman, N.G., 1991b. Age assessment of fallow deer (*Dama dama*): from a scoring scheme based on radiographs of developing permanent molariform teeth. *J. Zool.* 224, 367–379.
- Buchan, M., Müldner, G., Ervynck, A., Britton, K., 2016. Season of birth and sheep husbandry in late Roman and Medieval coastal Flanders: a pilot study using tooth enamel  $\delta^{18}\text{O}$  analysis. *Environ. Archaeol.* 21, 260–270. <http://dx.doi.org/10.1179/1749631414Y.00000000055>.
- Budd, P., Montgomery, J., Barreiro, B., Thomas, R.G., 2000. Differential diagenesis of strontium in archaeological human dental tissues. *Appl. Geochem.* 15, 687–694.
- Cameron, R.D., Reed, D.J., Dau, J.R., Smith, W.T., 1992. Redistribution of calving caribou in response to oil field development on the Arctic slope of Alaska. *Arctic* 45, 338–342.
- Capo, R.C., Stewart, B.W., Chadwick, O.A., 1998. Strontium isotopes as tracers of ecosystem processes: theory and methods. *Geoderma* 82, 197–225.
- Charlier, B.L.A., Ginibre, C., Morgan, D., Nowell, G.M., Pearson, D.G., Davidson, J.P., Otley, C.J., 2006. Methods for the microsampling and high-precision analysis of strontium and rubidium isotopes at single crystal scale for petrological and geochemical applications. *Chem. Geol.* 232, 114–133. <http://dx.doi.org/10.1016/j.chemgeo.2006.02.015>.
- Chisholm, B.S., Nelson, D.E., Schwarcz, H.P., 1982. Stable carbon isotope ratios as a measure of marine versus terrestrial protein in ancient diets. *Science* 216, 1131–1132.
- Copeland, S.R., Sponheimer, M., le Roux, P.J., Grimes, V., Lee-Thorp, J.A., de Ruiter, D.J., Richards, M.P., 2008. Strontium isotope ratios ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) of tooth enamel: a comparison of solution and laser ablation multicollector inductively coupled plasma mass spectrometry methods. *Rapid Commun. Mass Spectrom.* 22, 3187–3194. <http://dx.doi.org/10.1002/rcm.3717>.
- Couturier, S., Brunelle, J., Vandal, D., St-Martin, G., 1990. Changes in the population dynamics of the George River caribou herd 1976–87. *Arctic* 43, 9–20.
- Craig, O., Ross, R., Andersen, S.H., Milner, N., Bailey, G.N., 2006. Focus: sulphur isotope variation in archaeological marine fauna from northern Europe. *J. Archaeol. Sci.* 33, 1642–1646. <http://dx.doi.org/10.1016/j.jas.2006.05.006>.
- Dahle, B., Reimers, E., Colman, J.E., 2008. Reindeer (*Rangifer tarandus*) avoidance of a highway as revealed by lichen measurements. *Eur. J. Wildl. Res.* 54, 27–35. <http://dx.doi.org/10.1007/s10344-007-0103-5>.
- Dansgaard, W., 1964. Stable isotopes in precipitation. *Tellus* 16, 436–468. <http://dx.doi.org/10.3402/tellusa.v16i4.8993>.
- Deniel, C., Pin, C., 2001. Single-stage method for the simultaneous isolation of lead and strontium from silicate samples for isotopic measurements. *Anal. Chim. Acta* 426, 95–103. [http://dx.doi.org/10.1016/S0003-2670\(00\)01185-5](http://dx.doi.org/10.1016/S0003-2670(00)01185-5).
- DeNiro, M.J., Epstein, S., 1978. Influence of diet on the distribution of carbon isotopes in animals. *Geochim. Cosmochim. Acta* 42, 495–506.
- DeNiro, M.J., Epstein, S., 1981. Influence of diet on the distribution of nitrogen isotopes in animals. *Geochim. Cosmochim. Acta* 45, 341–351.
- D'Arrigo, R., Mashig, E., Frank, D., Wilson, R., Jacoby, G., 2005. Temperature variability over the past millennium inferred from Northwestern Alaska tree rings. *Clim. Dyn.* 24, 227–236.
- Domingo, L., Pérez-Dios, P., Fernández, M.H., Martín-Chivelet, J., Ortiz, J.E., Torres, T., 2015. Late Quaternary climatic and environmental conditions of northern Spain: an isotopic approach based on the mammalian record from La Paloma cave. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 440, 417–430. <http://dx.doi.org/10.1016/j.palaeo.2015.09.017>.
- Eckardt, H., Chenery, C., Booth, P., Evans, J.A., Lamb, A., Müldner, G., 2009. Oxygen and strontium isotope evidence for mobility in Roman Winchester. *J. Archaeol. Sci.* 36, 2816–2825. <http://dx.doi.org/10.1016/j.jas.2009.09.010>.
- Evans, J., Stoodley, N., Chenery, C., 2006. A strontium and oxygen isotope assessment of a possible fourth century immigrant population in a Hampshire cemetery, southern England. *J. Archaeol. Sci.* 33, 265–272. <http://dx.doi.org/10.1016/j.jas.2005.07.011>.
- Farrell, T.F., Jordan, P., Taché, K., Lucquin, A., Gibbs, K., Jorge, A., Britton, K., Craig, O.E., Knecht, R., 2014. Specialized processing of aquatic resources in prehistoric Alaskan pottery? A lipid-residue analysis of ceramic sherds from the Thule-period site of Nunalleq, Alaska. *Arct. Anthropol.* 51, 86–100. <http://dx.doi.org/10.3368/aa.51.1.86>.
- Feranec, R.S., MacFadden, B.J., 2000. Evolution of the grazing niche in Pleistocene mammals from Florida: evidence from stable isotopes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 162, 155–169.
- Feranec, R.S., Hadly, E.A., Paytan, A., 2007. Determining landscape use of Holocene mammals using strontium isotopes. *Oecologia* 153, 943–950. <http://dx.doi.org/10.1007/s00442-007-0779-y>.
- Feranec, R.S., Hadly, E.A., Paytan, A., 2009. Stable isotopes reveal seasonal competition for resources between late Pleistocene bison (*bison*) and horse (*equus*) from Rancho La Brea, Southern California. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 271, 153–160. <http://dx.doi.org/10.1016/j.palaeo.2008.10.005>.
- Ferguson, S.H., Elkie, P.C., 2004. Seasonal movement patterns of woodland caribou (*Rangifer tarandus caribou*). *J. Zool.* 262, 125–134. <http://dx.doi.org/10.1017/S0952836903004552>.
- Fienup-Riordan, A., 2000. *Hunting Tradition in a Changing World: Yup'ik Lives in Alaska Today*. Rutgers University Press, New Jersey.
- Fleming, M.D., Chapin, F.S., Cramer, W., Hufford, G.L., Serreze, M.C., 2000. Geographic patterns and dynamics of Alaskan climate interpolated from a sparse station record. *Glob. Chang. Biol.* 6, 49–58. <http://dx.doi.org/10.1046/j.1365-2486.2000.06008.x>.
- Forbes, V., Britton, K., Knecht, R., 2015. Preliminary archaeoentomological analyses of permafrost-preserved cultural layers from the pre-contact Yup'ik Eskimo site of Nunalleq, Alaska: implications, potential and methodological considerations. *Environ. Archaeol.* 20, 158–167. <http://dx.doi.org/10.1179/1749631414Y.0000000037>.
- Ford, J.D., Smit, B., Wandel, J., 2006. Vulnerability to climate change in the Arctic: a case study from Arctic Bay, Canada. *Glob. Environ. Chang.* 16, 145–160. <http://dx.doi.org/10.1016/j.gloenvcha.2005.11.007>.
- Fricke, H.C., Clyde, W.C., O'Neil, J.R., 1998. Intra-tooth variations in  $\delta^{18}\text{O}$  ( $\text{PO}_4$ ) of mammalian tooth enamel as a record of seasonal variations in continental climate variables. *Geochim. Cosmochim. Acta* 62, 1839–1850. [http://dx.doi.org/10.1016/S0016-7037\(98\)00114-8](http://dx.doi.org/10.1016/S0016-7037(98)00114-8).
- Fuller, B.T., Richards, M.P., Mays, S.A., 2003. Stable carbon and nitrogen isotope variations in tooth dentine serial sections from Wharram Percy. *J. Archaeol. Sci.* 30, 1673–1684. [http://dx.doi.org/10.1016/S0305-4403\(03\)00073-6](http://dx.doi.org/10.1016/S0305-4403(03)00073-6).
- Gat, J.R., 1980. The isotopes of hydrogen and oxygen in precipitation. In: Fritz, P., Fontes, J.C. (Eds.), *Handbook of Environmental Isotope Geochemistry*, Vol. 1: The Terrestrial Environment. Elsevier, Amsterdam.
- Gat, J.R., 1996. Oxygen and hydrogen isotopes in the hydrologic cycle. *Annu. Rev. Earth Planet. Sci.* 24, 225–262. <http://dx.doi.org/10.1146/annurev.earth.24.1.225>.
- Green, G.P., Bestland, E.A., Walker, G.S., 2004. Distinguishing sources of base cations in irrigated and natural soils: evidence from strontium isotopes. *Biogeochemistry* 68, 199–225.
- Gunn, A., 2003. Voles, lemmings and caribou-population cycles revisited? *Rangifer* 14, 105–111.
- Gunn, A., Miller, F.L., 1986. Traditional behaviour and fidelity to yup'ik calving grounds by barren-ground caribou. *Rangifer* 6, 151–158.
- Hansman, H., 2016. As climate change heats up, arctic residents struggle to keep their homes. *Guardian*. <https://www.theguardian.com/environment/2016/mar/15/climate-change-alaska-ice-melt-weather-resistant-housing> (15th March).
- Hedges, R.E., Stevens, R.E., Richards, M.P., 2004. Bone as a stable isotope archive for local climatic information. *Quat. Sci. Rev.* 23, 959–965. <http://dx.doi.org/10.1016/j.quascirev.2003.06.022>.
- Hegel, T.M., Russell, K., 2013. Status of northern mountain caribou (*Rangifer tarandus caribou*) in Yukon, Canada. *Rangifer* 33, 59–70.
- Hemming, J.E., 1971. *The Distribution and Movement Patterns of Caribou in Alaska*. Alaska Department of Fish and Game, pp. 1–60.
- Henton, E., Martin, L., Garrard, A., Jourdan, A.-L., Thirlwall, M., Boles, O., 2017. Gazelle seasonal mobility in the Jordanian steppe: the use of dental isotopes and microwear as environmental markers, applied to Epipalaeolithic Kharaneh IV. *J. Archaeol. Sci.* Rep. 11, 147–158. <http://dx.doi.org/10.1016/j.jasrep.2016.11.031>.
- Hinzman, L.D., Bettz, N.D., Bolton, W.R., Chapin, F.S., Dyrurgorov, M.B., Fastie, C.L., Griffith, B., Hollister, R.D., Hope, A., Huntington, H.P., Jensen, A.M., Jia, G.J., Jorgenson, T., Kane, D.L., Klein, D.R., Kofinas, G., Lynch, A.H., Lloyd, A.H., McGuire, A.D., Nelson, F.E., Oechel, W.C., Osterkamp, T.E., Racine, C.H., Romanovsky, V.E., Stone, R.S., Stow, D.A., Sturm, M., Tweedie, C.E., Vourlitis, G.L., Walker, M.D., Walker, D.A., Webber, P.J., Welker, J.M., Winker, K.S., Yoshikawa, K., 2005. Evidence and implications of recent climate change in northern Alaska and other Arctic regions. *Clim. Chang.* 72, 251–298. <http://dx.doi.org/10.1007/s10584-005-5352-2>.
- Hobson, K.A., 1999. Tracing origins and migration of wildlife using stable isotopes: a review. *Oecologia* 120, 314–326.
- Hoppe, K.A., 2004. Late Pleistocene mammoth herd structure, migration patterns, and Clovis hunting strategies inferred from isotopic analyses of multiple death assemblages. *Paleobiology* 30, 129–145. [http://dx.doi.org/10.1666/0094-8373\(2004\)030<0129:LPMHSM>2.0.CO;2](http://dx.doi.org/10.1666/0094-8373(2004)030<0129:LPMHSM>2.0.CO;2).
- Hoppe, K.A., Koch, P.L., Carlson, R.W., Webb, S.D., 1999. Tracking mammoths and mastodons: reconstruction of migratory behaviour using strontium isotope ratios. *Geology* 27, 439–442. [http://dx.doi.org/10.1130/0091-7613\(1999\)027<0439:TMAMRO>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1999)027<0439:TMAMRO>2.3.CO;2).
- Jacobs, J.D., Maarouf, A.R., Perkins, E.A., 1996. The recent record of climate on the range of the George River Caribou Herd, Northern Québec and Labrador, Canada. *Rangifer* 16, 193–200.
- Jakimchuk, R.D., Ferguson, S.H., Sopuck, L.G., 1987. Differential habitat use and sexual segregation in the Central Arctic caribou herd. *Can. J. Zool.* 65, 534–541. <http://dx.doi.org/10.1139/z87-083>.
- Johnson, C.M., Fridrich, C.J., 1990. Non-monotonic chemical and O, Sr, Nd, and Pb isotope zonations and heterogeneity in the mafic to silicic-composition magma chamber of the Grizzly Peak Tuff, Colorado. *Contrib. Mineral. Petrol.* 105, 677–690. <http://dx.doi.org/10.1007/BF00306533>.
- Julien, M.-A., Bocherens, H., Burke, A., Drucker, D.G., Patou-Mathis, M., Krotova, O., Péan, S., 2012. Were European steppe bison migratory?  $^{18}\text{O}$ ,  $^{13}\text{C}$  and Sr intra-tooth isotopic variations applied to a palaeoethological reconstruction. *Quat. Int.* 271, 106–119. <http://dx.doi.org/10.1016/j.quaint.2012.06.011>.
- Koch, P.L., Tuross, N., Fogel, M.L., 1997. The effects of sample treatment and diagenesis on the isotopic integrity of carbonate in biogenic hydroxylapatite. *J. Archaeol. Sci.* 24, 417–429.
- Kohn, M.J., 1996. Predicting animal  $\delta^{18}\text{O}$ : accounting for diet and physiological adaptation. *Geochim. Cosmochim. Acta* 60, 4811–4829.
- Kohn, M.J., 2004. Comment: tooth enamel mineralization in ungulates: implications for recovering a primary isotopic time-series, by B.H. Passey and T.E. Cerling (2002). *Geochim. Cosmochim. Acta* 68, 403–405. [http://dx.doi.org/10.1016/S0016-7037\(03\)00443-5](http://dx.doi.org/10.1016/S0016-7037(03)00443-5).
- Kohn, M.J., Cerling, T.E., 2002. Stable isotope compositions of biological apatite. *Rev. Mineral. Geochem.* 48, 455–488. <http://dx.doi.org/10.2138/rmg.2002.48.12>.
- Kohn, M., Schoeninger, M., Valley, J., 1996. Herbivore tooth oxygen isotope



- compositions: effects of diet and physiology. *Geochim. Cosmochim. Acta* 60, 3889–3896.
- Köppen, W., 1900. Versuch einer Klassifikation der Klimate, vorzugsweise nach ihren Beziehungen zur Pflanzenwelt. *Geogr. Z.* 6, 593–611.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World map of the Köppen-Geiger climate classification updated. *Meteorol. Z.* 15, 259–263. <http://dx.doi.org/10.1127/0941-2948/2006/0130>.
- Ledger, P.M., Forbes, V., Masson-MacLean, E., Knecht, R.A., 2016. Dating and digging stratified archaeology in circumpolar North America: a view from Nunalleq, south-western Alaska. *Arctic* 69, 378. <http://dx.doi.org/10.14430/arctic4599>.
- Longinelli, A., 1984. Oxygen isotopes in mammal bone phosphate: a new tool for paleohydrological and paleoclimatological research? *Geochim. Cosmochim. Acta* 48, 385–390.
- Luz, B., Kolodny, Y., Horowitz, M., 1984. Fractionation of oxygen isotopes between mammalian bone-phosphate and environmental drinking water. *Geochim. Cosmochim. Acta* 48, 1689–1693. [http://dx.doi.org/10.1016/0016-7037\(84\)90338-7](http://dx.doi.org/10.1016/0016-7037(84)90338-7).
- Mann, M.E., 2002. Little Ice Age. In: Munn, R.E. (Ed.), *Encyclopaedia of Global Environmental Change*. Wiley, Chichester; New York, pp. 504–509.
- Mann, M.E., Bradley, R.S., Hughes, M.K., 1999. Northern hemisphere temperatures during the past millennium: inferences, uncertainties, and limitations. *Geophys. Res. Lett.* 26, 759–762. <http://dx.doi.org/10.1029/1999GL900070>.
- Masood, S., Van Zuiden, T.M., Rodgers, A.R., Sharma, S., 2017. An uncertain future for woodland caribou (*Rangifer tarandus caribou*): the impact of climate change on winter distribution in Ontario. *Rangifer* 37, 11–30. <http://dx.doi.org/10.7557/2.37.1.4103>.
- Maurer, A.-F., Galer, S.J.G., Knipper, C., Beierlein, L., Nunn, E.V., Peters, D., Tütken, T., Alt, K.W., Schöne, B.R., 2012. Bioavailable  $^{87}\text{Sr}/^{86}\text{Sr}$  in different environmental samples-effects of anthropogenic contamination and implications for isoscapes in past migration studies. *Sci. Total Environ.* 433, 216–229.
- McConnell, D., 1973. *Apatite, Its Crystal Chemistry, Mineralogy, Utilization and Geologic and Biologic Occurrences*. Springer-Verlag, Wien.
- McManus-Fry, E., 2015. Pre-contact Ecology, Subsistence, and Diet on the Yukon Kuskokwim Delta: An Integrated Ecosystem Approach to Pre-contact Arctic Lifeways Using Zooarchaeological Analysis and Stable Isotope Techniques. PhD Thesis. University of Aberdeen.
- McManus-Fry, E., Knecht, R., Dobney, K., Richards, M.P., Britton, K., 2016. Dog-human dietary relationships in Yup'ik western Alaska: the stable isotope and zooarchaeological evidence from pre-contact Nunalleq. *J. Archaeol. Sci. Rep.* <http://dx.doi.org/10.1016/j.jasrep.2016.04.007>.
- McNeil, P., Russell, D.E., Griffith, B., Gunn, A., Kofinas, G.P., 2005. Where the wild things are: seasonal variation in caribou distribution in relation to climate change. *Rangifer* 16, 51–63.
- Meldgaard, M., 1986. The Greenland Caribou- zoogeography, taxonomy, and population dynamics, Meddelelser om Grønland. Bioscience 20, 1–88 (Copenhagen).
- Metcalfe, J.Z., Longstaffe, F.J., 2014. Environmental change and seasonal behaviour of mastodons in the Great Lakes region inferred from stable isotope analysis. *Quat. Res.* 82, 366–377. <http://dx.doi.org/10.1016/j.yqres.2014.07.002>.
- Montgomery, J., Evans, J.A., Cooper, R.E., 2007. Resolving archaeological populations with Sr-isotope mixing models. *Appl. Geochem.* 22, 1502–1514.
- Murie, O.J., 1935. Alaska-Yukon caribou. *North American Fauna* 54, 1–92.
- Passey, B.H., Cerling, T.E., 2002. Tooth enamel mineralization in ungulates: implications for recovering a primary isotopic time-series. *Geochim. Cosmochim. Acta* 66, 3225–3234.
- Pellegrini, M., Donahue, R.E., Chenery, C., Evans, J., Lee-Thorp, J., Montgomery, J., Mussi, M., 2008. Faunal migration in late-glacial central Italy: implications for human resource exploitation. *Rapid Commun. Mass Spectrom.* 22, 1714–1726. <http://dx.doi.org/10.1002/rcm.3521>.
- Pfister, C., 1995. Monthly temperature and precipitation in central Europe 1525–1979: quantifying documentary evidence on weather and its effects. In: Bradley, R.S., Jones, P.D. (Eds.), *Climate Since AD 1500*. Routledge, London.
- Pilkington, E., 2008. *The Village at the Tip of the Iceberg*. The Guardian.
- Post, E., Forchhammer, M.C., Bret-Harte, M.S., Callaghan, T.V., Christensen, T.R., Elberling, B., Fox, A.D., Gilg, O., Hik, D.S., Hoye, T.T., Ims, R.A., Jeppesen, E., Klein, D.R., Madsen, J., McGuire, A.D., Rysgaard, S., Schindler, D.E., Stirling, I., Tamstorf, M.P., Tyler, N.J.C., van der Wal, R., Welker, J., Wookey, P.A., Schmidt, N.M., Aastrup, P., 2009. Ecological dynamics across the Arctic associated with recent climate change. *Science* 325, 1355–1358. <http://dx.doi.org/10.1126/science.1173113>.
- Price, T.D., Manzanilla, L., Middleton, W.D., 2000. Immigration and the ancient city of Teotihuacan in Mexico: a study using strontium isotope ratios in human bone and teeth. *J. Archaeol. Sci.* 27, 903–913. <http://dx.doi.org/10.1006/jasc.1999.0504>.
- Price, T.D., Burton, J.H., Bentley, R.A., 2002. The characterization of biologically available strontium isotope ratios for the study of prehistoric migration. *Archaeometry* 44, 117–135.
- Price, T.D., Meiggs, D., Weber, M.J., Pike-Tay, A., 2017. The migration of late Pleistocene reindeer: isotopic evidence from northern Europe. *Archaeol. Anthropol. Sci.* 9, 371–394. <http://dx.doi.org/10.1007/s12520-015-0290-z>.
- Root, T.L., Price, J.T., Hall, K.R., Schneider, S.H., Rosenzweig, C., Pounds, J.A., 2003. Fingerprints of global warming on wild animals and plants. *Nature* 421, 57–60. <http://dx.doi.org/10.1038/nature01333>.
- Russell, D.E., Martell, A.M., Nixon, W.A.C., 1993. Range ecology of the porcupine caribou herd in Canada. *Rangifer* 13, 1–168.
- Ruttan, R.A., 2012. New caribou crisis- then and now. *Rangifer* 20, 85–102.
- Sayle, K.L., Hamilton, D.H., Cook, G.T., Ascough, P.L., Gestsdottir, H., McGovern, T.H., 2016. Deciphering diet and monitoring movement: multiple stable isotope analysis of the Viking age settlement at Hofstaðir, Lake Myvatn, Iceland. *Am. J. Phys. Anthropol.* 160, 126–136. <http://dx.doi.org/10.1002/ajpa.22939>.
- Schoeninger, M.J., DeNiro, M.J., Tauber, H., 1983. Stable nitrogen isotope ratios of bone collagen reflect marine and terrestrial components of prehistoric human diet. *Science* 220, 1381–1383.
- Simon, J., 2016. Customary and Traditional Use Worksheet, Caribou, Game Management Unit 18. Alaska Department of Fish and Game, Division of Subsistence (Special Publication No. BOG 2016-03, Fairbanks).
- Simon-Sakurai, K.A., 2013. *Never Alone: Narratives of Spirits in an Alaskan Yup'ik Community*. PhD Thesis. University of Aberdeen.
- Skoog, R.O., 1968. *Ecology of the Caribou (Rangifer tarandus granti) in Alaska*. PhD Thesis. University of California (Berkeley).
- Stevens, R.E., Hermoso-Buxan, X.L., Marin-Arroyo, A.B., Gonzalez-Morales, M.R., Straus, L.G., 2014. Investigation of late Pleistocene and early Holocene palaeoenvironmental change at El Miron cave (Cantabria, Spain): insights from carbon and nitrogen isotope analyses of red deer. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 414, 46–60. <http://dx.doi.org/10.1016/j.palaeo.2014.05.049>.
- Tape, K.D., Gustine, D.D., Ruess, R.W., Adams, L.G., Clark, J.A., 2016. Range expansion of moose in arctic Alaska linked to warming and increased shrub habitat. *PLoS One* 11, e0152636. <http://dx.doi.org/10.1371/journal.pone.0152636>.
- Tauber, H., 1981.  $^{13}\text{C}$  evidence for dietary habits of prehistoric man in Denmark. *Nature* 292, 332–333.
- Tornero, C., Balasse, M., Bălăşescu, A., Chataigner, C., Gasparyan, B., Montoya, C., 2016. The altitudinal mobility of wild sheep at the Epigravettian site of Kalavan 1 (Lesser Caucasus, Armenia): evidence from a sequential isotopic analysis in tooth enamel. *J. Hum. Evol.* 97, 27–36. <http://dx.doi.org/10.1016/j.jhevol.2016.05.001>.
- Tornero, C., Aguilera, M., Ferrio, J.P., Arcusa, H., Moreno-García, M., García-Reig, S., Rojo-Guerra, M., 2017. Vertical sheep mobility along the altitudinal gradient through stable isotope analyses in tooth molar bioapatite, meteoric water, and pastures: a reference from the Ebro valley to the Central Pyrenees. *Quat. Int.* <http://dx.doi.org/10.1016/j.quaint.2016.11.042>.
- Tütken, T., Langenegger, E., Wild, W., 2008. Einheimisch oder fremd? Isotopenanalyse einer Frau des 9. Jh. n. Chr. aus Elsau, Kanton Zürich, Schweiz. *Anthropol. Anz.* 66, 1–32.
- Valenzuela-Lamas, S., Jiménez-Manchón, S., Evans, J., López, D., Jornet, R., Albarella, U., 2016. Analysis of seasonal mobility of sheep in Iron Age Catalonia (north-eastern Spain) based on strontium and oxygen isotope analysis from tooth enamel: first results. *J. Archaeol. Sci. Rep.* 6, 828–836. <http://dx.doi.org/10.1016/j.jasrep.2015.08.042>.
- Vitousek, P.M., Kennedy, M.J., Derry, L.A., Chadwick, O.A., 1999. Weathering versus atmospheric sources of strontium in ecosystems on young volcanic soils. *Oecologia* 121, 255–259.
- Weladji, R.B., Klein, D.R., Holand, Ø., Mysterud, A., 2002. Comparative response of *Rangifer tarandus* and other northern ungulates to climatic variability. *Rangifer* 22, 33–50.
- Whipkey, C.E., Capo, R.C., Chadwick, O.A., Stewart, B.W., 2000. The importance of sea spray to the cation budget of a coastal Hawaiian soil: a strontium isotope approach. *Chem. Geol.* 168, 37–48. [http://dx.doi.org/10.1016/S0009-2541\(00\)00187-X](http://dx.doi.org/10.1016/S0009-2541(00)00187-X).
- Widga, C., Walker, J.D., Stockli, L.D., 2010. Middle Holocene Bison diet and mobility in the eastern Great Plains (USA) based on  $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$ , and  $^{87}\text{Sr}/^{86}\text{Sr}$  analyses of tooth enamel carbonate. *Quat. Res.* 73, 449–463. <http://dx.doi.org/10.1016/j.yqres.2009.12.001>.
- Wißing, C., Matzerath, S., Turner, E., Bocherens, H., 2015. Paleoclimatic and climatic implications of stable isotope results from late Pleistocene bone collagen, Ziegeleigrube Coenen, Germany. *Quat. Res.* 84, 96–105. <http://dx.doi.org/10.1016/j.yqres.2015.05.005>.
- Zazzo, A., Balasse, M., Patterson, W., 2006. The reconstruction of mammal individual history: refining high-resolution isotope record in bovine tooth dentine. *J. Archaeol. Sci.* 33, 1177–1187 doi: 10.1016/j.jas.2005.12.006.
- Zazzo, A., Bendrey, R., Vella, D., Moloney, A.P., Monahan, F.J., Schmidt, O., 2012. A refined sampling strategy for intra-tooth stable isotope analysis of mammalian enamel. *Geochim. Cosmochim. Acta* 84, 1–13. <http://dx.doi.org/10.1016/j.gca.2012.01.012>.